www.dynamicpapers.com



Mark Scheme (Results)

October 2020

Pearson Edexcel IAL In Statistics 1
Paper WST01/01

www.dynamicpapers.com

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

October 2020
Publications Code WST01_01_2010_MS
All the material in this publication is copyright
© Pearson Education Ltd 2020

www.dynamicpapers.com

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

www.dynamicpapers.com

EDEXCEL IAL MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 75.

2. The Edexcel Mathematics mark schemes use the following types of marks:

- **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
- **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
- **B** marks are unconditional accuracy marks (independent of M marks)
- Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
- ft – follow through
- the symbol $\surd$ will be used for correct ft
- cao – correct answer only
- cso - correct solution only. There must be no errors in this part of the question to obtain this mark
- isw – ignore subsequent working
- awrt – answers which round to
- SC: special case
- oe – or equivalent (and appropriate)
- dep – dependent
- indep – independent
- dp decimal places
- sf significant figures
- * The answer is printed on the paper
- ⌐ The second mark is dependent on gaining the first mark

4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.

6. If a candidate makes more than one attempt at any question:
   - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
   - If either all attempts are crossed out or none are crossed out, mark the last most complete solution.

7. Ignore wrong working or incorrect statements following a correct answe

www.dynamicpapers.com

Question Number	Scheme	Marks
1.	x | -1 | 2 | 3 | 4 | 7 $\mathrm{P}(X=x)$ | $\frac{9}{k}$ | $\frac{6}{k}$ | $\frac{5}{k}$ | $\frac{4}{k}$ | $\frac{1}{k}$	M1
	$\sum \mathrm{P}(X=x)=1 \Rightarrow \quad \frac{25}{k}=1$	M1
	$k=25$	A1
	$\mathrm{E}(X)=\frac{1}{25}\left[-1\times 9+2\times 6+3\times 5+4\times 4+7\times 1\right]$	M1
	$=\underline{\underline{\frac{41}{25}}}$	A1
		[5]
	Notes	
	1st M1 for at least 3 correct probabilities in terms of k (may be seen used in expression for E(X)) 2nd M1 for attempting to use sum of 5 probs = 1 (ft their probabilities) 1st A1 for $k = 25$ (stated or used correctly) 3rd M1 for attempt at a correct expression at least 3 products (ft their k – value or letter) 2nd A1 for $\frac{41}{25}$ or exact equivalent e.g. 1.64 Correct answer with no incorrect method marks scores 5/5	

www.dynamicpapers.com

Question Number	Scheme	Marks
2. (a)		B1 B1 (2)
(b)	P(W) = 0.4p + 0.35q + "0.25"×0.4 [= 0.4p + 0.35q + 0.1]	B1ft (1)
(c)	Correct expression: $P(W \cap V) = "0.1" = "0.25" \times P(W)$ or P(W) = P($W \mid V$) = 0.4 0.1 = 0.25(0.4p + 0.35q + 0.25×0.4) or 0.4p + 0.35q + 0.25×0.4 = 0.4	M1 A1 (2)
(d)	$\frac{7}{30} = \frac{0.35(1-q)}{"P(J)"}$ Since V and W are independent so are V and $W' = J$ so P(J) = 0.6 or sub P(J) = 1 – their (b) to get an equation in p and q [May see 8p – 23q +12 = 0] [So $1-q = \frac{2}{3}P(J)$ therefore] ***q* = 0.6** $8p + 7 \times "0.6" = 6$ So ***p* = 0.225** or $\frac{9}{40}$	M1 dM1 A1 ddM1 A1 (5)
(e)	{P($V \mid W$) = P(V) = 0.25 (since independent) and P($M \mid W$) = 0.225 (= p)} P($F \mid W$) = $\frac{0.35 \times "0.6"}{"0.4"}$ or $\frac{0.35q}{(b)}$; = $\frac{21}{40}$ or 0.525 [Since this prob > 0.5 therefore it must be the largest] so conclusion is correct Allow B1ft for comparing 3 calculated probs of the form P($M \cap W$) needn't be correct ft	M1 ;A1 B1ft (3) **[13]**

	Notes
(a)	1st B1 0.25 for P(V) 2nd B1 for correct probabilities on 2nd branches (1 – p), (1 – q) [allow their values] and 0.6
(b)	B1ft for a correct expression using their values from tree diagram
(c)	M1 for sight or use of a correct expression in V and W or correct equation in p and q (ft their part (b)) A1 for a fully correct equation (needn't be simplified) [may see 0.4p + 0.35q = 0.3 or 8p + 7q = 6]
(d)	1st M1 for using given conditional probability to form an equation in q and P(J) using $\frac{7}{30}$ 2nd dM1 (dep on 1st M1) for a getting P(J) = 0.6 or sub 1 – their (b) and get 2nd equation in p and q 1st A1 for q = 0.6 [NB must be q = 0.6 not just P(J) = 0.6] May see after 3rd M1 for solving with p 3rd ddM1(dep on both Ms) for seeing substitution of their 1st value to find the 2nd value (p or q) Allow ft of their p or q in one of their equations provided p and q both lie in (0, 1) 2nd A1 for p = 0.225 or exact equivalent After the 2nd M1, sight of p = 0.225 and q = 0.6 earns the final 3 marks
(e)	M1 for a method for finding P($F \mid W$) A1 for a correct value $\frac{21}{40}$ or exact equivalent B1ft for a correct conclusion based on enough probs found ft their probabilities

www.dynamicpapers.com

Question Number	Scheme	Marks
3. (a)	[D = distance achieved] $\mathrm{P}(D > 4.3) = \mathrm{P}\left(Z > \dfrac{4.3-3.8}{0.9}\right)$ or $\mathrm{P}(Z > 0.555\ldots)$	M1
	$= 1 - 0.7123$ (tables)	M1
	$= 0.2877$ (tables) or $0.289257\ldots$ (calc) awrt **0.288** or awrt **0.289**	A1
		(3)
(b)	$\dfrac{d-3.8}{0.9} = -0.8416$ (calc $-0.84162123\ldots$)	M1 ; B1
	$d = 3.0425\ldots$ awrt **3.04**	A1
		(3)
(c)	$\mathrm{P}(D > g \mid D > 4.3) = \dfrac{\mathrm{P}(D > g)}{\mathrm{P}(D > 4.3) \text{ or (a)}}$ $\left[= \dfrac{1}{3}\right]$ (o.e.)	M1
	$\therefore \mathrm{P}(D > g) = \frac{1}{3}(\text{a}) = 0.096419\ldots$	A1ft (o.e)
	$\dfrac{g-3.8}{0.9} = 1.302228\ldots$	dM1
	so $g = 4.97200..$ awrt **4.97** or awrt **4.98**	A1
		(4)
(d)	P(no gold medals) $= \left(\frac{2}{3}\right)^3$	M1
	P(at least one gold) $= 1 - \left(\frac{2}{3}\right)^3$	M1
	$= \frac{19}{27}$	A1
		(3)
		[13]

	Notes
(a)	1st M1 for standardising 4.3 with 3.8 and 0.9 (allow $\pm$) 2nd M1 for $1 - p$ (where $0.7 < p < 0.8$) A1 for awrt 0.288 or 0.289 (calc. 0.289257) (correct answer only 3/3)
(b)	M1 for standardising with d, 3.8 and 0.9 and setting equal to a z value $0.8 < \lvert z \rvert < 0.9$ B1 for $z = \pm 0.8416$ or better used A1 for awrt 3.04 (condone $d \geqslant \ldots$)
Ans only	For awrt 3.0425 or 3.0426 score 3/3 For awrt 3.04 score M1B0A1
(c)	1st M1 for either expression for the conditional prob. [or sight of $\frac{1}{3}$(a)] (ft their answer to (a) to 2 sf) 1st A1ft for $\mathrm{P}(D > g) = 0.096$ or better (0.289 gives 0.09633… calc 0.096419…) The $\mathrm{P}(D > g)$ may be clearly shown on a diagram. 1st M1A1 can be awarded for $\mathrm{P}(D > g) = \frac{1}{3}$(a) or for $\mathrm{P}(D < g) = 1 - \frac{1}{3}$(a) [ft their (a) to 2 sf] 2nd dM1 (dep on 1st M1) for standardising with g, 3.8 and 0.9 and put equal to a z value where $\lvert z \rvert > 1$ 2nd A1 for awrt 4.97 or 4.98 (Correct answer with no incorrect working seen 4/4) (condone $g \geqslant \ldots$)
SC	**(Medals v Certificates)** 1st B1 for $[\mathrm{P}(D > g) =] \frac{1}{3} \times 0.8 = \frac{4}{15}$ or 0.267 (score as 1st M0 1st A1) 2nd B1 for g = awrt 4.36 (4.358 tables, 4.3606..calc) (score as 2nd M0 2nd A1)
(d)	1st M1 for a correct probability of no gold medals or 2 of: $3\left(\frac{2}{3}\right)^2 \times \frac{1}{3}$ or $3\left(\frac{1}{3}\right)^2 \times \frac{2}{3}$ or $\left(\frac{1}{3}\right)^3$ 2nd M1 for $1 - p^3$ or $3(p)^2(1-p) + 3p(1-p)^2 + (1-p)^3$ where $0 < p < 1$ A1 for $\frac{19}{27}$ (or exact equivalent) only e.g. $0.\dot{7}0\dot{3}$

www.dynamicpapers.com

Question Number	Scheme	Marks
4. (a)	Upper quartile = 34 Lower limit = 24 – 15 = 9 or upper limit is "34" + 15 = 49 So outliers are: 8, 52.5 and 56	B1 M1 A1ft, A1ft (4)
(b)		B1 B1 B1 (3)
(c)	$Q_2 - Q_1 (=6) > ("4" =) Q_3 - Q_2$ or e.g. in words e.g. " Q_3 closer to Q_2 than Q_1 is" So negative (skew)	M1 A1ft (2)
(d)	IQR now "34" – 26 = 8 so new outlier limits are 26 – 1.5×"8" = **14** and "34" + 1.5×"8" = **46** 	M1 A1ft A1 (3)
(e)	[Q_1 has increased so both above 24 Median same so either side of or on median] So one **between 26 and 30** inc [Q_3 unchanged so must be either side of Q_3] so one **between "34" and 45** inc	B1 B1 (2) **[14]**

	Notes
(a)	B1 for Q_3 = 34 either stated or used/implied (score if seen on box plot) M1 for one correct calculation (ft their 34 for upper limit) [May be implied by correct outliers] 2nd A1ft for the lower **outlier** at 8 (ft their limit provided limit ⩽ 12) 3rd A1ft for upper **outliers** at 52.5 and 56 (ft their limit provided it is > 45) Award if their outliers are seen on box plot **NB These accuracy marks are for the outliers not the limits**
(b)	1st B1 for a box with Q_1 = 24, Q_2 = 30 Q_3 = their 34 and two whiskers one on each side. 2nd B1 for one lower whisker ending at 10 (or their 9) and outlier at 8 only 3rd B1 for one upper whisker ending at 45 (or their 49 to match "9") and outliers at 52.5 and 56 only
SC	**Extra whiskers. If one set of whiskers gives a correct box plot award B1B0B0** **Usual accuracy for plots – to within 0.5 of a square.**
(c)	M1 for correct comparison of $Q_2 - Q_1$ and $Q_3 - Q_2$ (ft their Q_3) (if no values seen must see comparison otherwise accept correctly assigned 6 and 4 without >) A1ft for correct deduction based on their Q_3 (+ve (skew) if their Q_3 > 36, no skew if their Q_3=36)
(d)	M1 for recognising new IQR and at least one correct new limit (ft their 34, implied by correct plot) 1st A1ft for a correct lower whisker ending at 15.5 (or their 14) and 2 correct outliers at 8 and 10 2nd A1 for a fully correct box plot with upper whisker to 45 (or could go to 46 [to match their 14])
SC	**Extra whiskers. If one set of whiskers gives a correct box plot award M1A0A1**
(e)	1st B1 for a range [26, 30] allow that (…) (o.e. eg between 26 and 30) 2nd B1 for a range [34, 45) condone […] or (…) (ft their 34 and allow o.e. e.g. between 34 and 45)

www.dynamicpapers.com

Question Number	Scheme	Marks
5. (a)	$y = 6.066 + 0.136\times 80$ $= 16.946$ (so annual rent is) **$ 16 946**	M1 A1 (2)
(b)	$S_{yy} = 3434 - \frac{183^2}{10}$ or $S_{xx} = 84\,818 - \frac{900^2}{10}$ $S_{yy} =$ **85.1** $S_{xx} =$ **3818**	M1 A1 A1 (3)
(c)	Need S_{xy} so use b so $S_{xy} = b\times S_{xx} = 0.136\times 3818$ or 519.248 $[r =] \frac{0.136\times "3818"}{\sqrt{"3818"\times "85.1"}}$ $= 0.9109448\ldots$ awrt **0.911**	M1; A1 M1 A1 (4)
(d)	Since (new $x = 90$ and [original or] new $\bar{x} = 90$) the term $(x - \bar{x})$ will be 0 Therefore (the 11th shop makes no change) S_{xy} stays the same	M1 A1 (2)
(e)	S_{xx} will be the same so b will be the same New $\bar{y} = \frac{183+15}{11} = 18$ (or a is reduced by 0.3) Equation is ***y* = 5.766 + 0.136*x***	M1 M1 A1 (3)
(f)	$x = 300$ is outside the range $300 \gg 90$ [$300 \gg 90 + 3\sigma = 90 + 3\times 18.63.. \approx 146$] So not suitable (since involves extrapolation) (o.e.)	B1 (1) **[15]**

	Notes
(a)	M1 for substituting $x = 80$ into the given equation A1 for awrt \$ 16 900 (or better)(allow "16.9 thousand dollars"). Must have some units. Condone $y =$
(b)	M1 for a correct expression for either (can be implied by sight of either correct answer) 1st A1 for 85.1 2nd A1 for 3818 or accept 3820
(c)	1st M1 for an attempt to use gradient of regression line to find S_{xy} 1st A1 for awrt 519 2nd M1 for a correct expression using their values (M0 if $S_{xy} = 900\times 183 = 164700$) 2nd A1 for awrt 0.911
(d)	M1 for stating or showing [old or] new $\bar{x}$ =90 (new $x = 90$ implied) or stating that $(x - \bar{x})$ term = 0 A1 for a fully correct argument mentioning new $x = \bar{x} = 90$ and that extra $(x - \bar{x})$ term = 0 Condone using $\bar{y} = 18.3$ instead of 18
(e)	1st M1 for a correct statement about S_{xx} or b (may be implied by 0.136 used correctly) 2nd M1 for a correct value for new $\bar{y}$ (calculation may be seen in (d) scores here when 18 is used) A1 for $y = 5.766$ (or awrt 5.77 or awrt 5.76) + $0.136x$ (correct equation scores 3/3)
(f)	B1 for suitable comparison (must see 300 vs 90 or 3000 vs 900) that says or implies that 300 will be outside the range and therefore not suitable. Not sufficient to just say "larger"

www.dynamicpapers.com

Question Number	Scheme	Marks
6. (a)	[E(A) =] $1\times0.4 + 4\times0.2 + 5\times0.25 + 7\times0.15$	M1
	$= \underline{\mathbf{3.5}}$ **(*)**	A1cso
		(2)
(b)	$\left[\mathrm{E}(A^2)=\right]1\times0.4 + 4^2\times0.2 + 5^2\times0.25 + 7^2\times0.15$ [= 17.2]	M1
	Var(A) = E(A^2) – [E(A)]^2 = 17.2 – 3.5^2	M1
	$= \underline{\mathbf{4.95}}$	A1
		(3)
(c)	(Discrete) uniform (distribution)	B1
		(1)
(d)	By symmetry $k = 6$	B1
		(1)
(e)	$\left[\text{Sam has } Z=\frac{3.5-4}{3}=-\frac{1}{6} \text{ and}\right]$ Tim needs $\frac{3.5-A}{4}<-\frac{1}{6}$ so $A > 4.166..$	M1
	So prob = 0.25 + 0.15 = $\underline{\mathbf{0.4}}$	A1
		(2)
(f)	Need largest possible $\mu = 7$ and smallest possible $\sigma = 1$	B1, B1
	P($X > 3.5$) is then $\mathrm{P}\left(Z>\frac{3.5-7}{1}\right)$ = P($Z > -3.5$)	M1
	= $\underline{\mathbf{0.9998}}$ (tables) or 0.999767…(calc)	A1
		(4)
(g)	[Need $A = 7$ and $B = 1$ (or ft from (f)) so] P($A = 7$)×P($B = 1$) <u>or</u> "0.15"×0.25	M1
	= $\underline{\mathbf{0.0375}}$	A1cso
		(2)
		[15]

	Notes
(a)	M1 for an attempt at E(A) – at least 3 correct products seen
	A1cso for 3.5 or exact equivalent with no incorrect working seen and M1 scored
(b)	1st M1 for an attempt at E(A^2) – at least 3 correct products
	2nd M1 for use of E(A^2) – [E(A)]^2 ft their value for E(A^2)
ALT	**M2** for $(-2.5)^2\times0.4 + (0.5)^2\times0.2 + (1.5)^2\times0.25 + (3.5)^2\times0.15$ (at least 3 correct products)
	A1 for 4.95 or an exact equivalent e.g. $\frac{99}{20}$
(c)	B1 for uniform (continuous uniform is B0)
(d)	B1 for stating $k = 6$ with a suitable reason e.g. mention of symmetry or full calculation
(e)	M1 for a suitable calculation for A e.g. $\frac{3.5-A}{4}<-\frac{1}{6}$ <u>or</u> stating $A = 5$ or 7 <u>or</u> $A >$ awrt 4.2 (o.e.)
	A1 for 0.4 (must be based on some correct calculation seen)
(f)	1st B1 for $\mu = 7$ may be implied from a standardisation with 3.5 seen
	2nd B1 for $\sigma = 1$ may be implied from a standardisation with 3.5 seen
	M1 for attempting correct probability i.e. P(Z … or X…) ft standardisation using 3.5, their $\mu \neq 4$ and their $\sigma \neq 3$ but their μ and σ must be "possible" values <u>or</u> P($Z > -3.5$)
	A1 for 0.9998 or better
(g)	M1 for "0.15"×0.25 ft their value of A from (f)
	A1cso for 0.0375 or exact equivalent e.g. $\frac{3}{80}$ (Must clearly come from $A = 7$ and $B = 1$ in (f))

G. B. Attwood 2/11/20

www.dynamicpapers.com

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom