www.dynamicpapers.com

Pearson Edexcel

Mark Scheme (Results)

January 2020

Pearson Edexcel In ational GCSE
In Further Pure Mathe ics (4PM1)
Paper 02R

www.dynamicpapers.com

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, ev ere

Pearson aspires to be the world's leading l company. Our aim is to help everyone progress in their lives through education. We bel every kind of learning, for all kinds of people, wherever they are in the world. We've been ed in education for over 150 years, and by working across 70 countries, in 100 langua e have built an international reputation for our commitment to high standards a ising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2020
Publications Code 4PM1_02R_2001_MS
All the material in this publication is copyright
© Pearson Education Ltd 2020

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme.
  Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners ar n doubt regarding the application of the mark scheme to a candidate's respon team leader must be consulted.
- Crossed out work sho marked UNLESS the candidate has replaced it with an alternative response.

- **Types of mark**
  - M marks: method marks
  - A marks: accuracy marks – c y be awarded when relevant M marks have been gained
  - B marks: unconditional accuracy m dependent of M marks)

- **Abbreviations**
  - cao – correct answer only
  - cso – correct solution only
  - ft – follow through
  - isw – ignore subsequent working
  - SC - special case
  - oe – or equivalent (and appropriate)
  - dep – dependent
  - indep – independent
  - awrt – answer which rounds to
  - eeoo – each error or omission

- **No working**
  If no working is shown then correct answers may score full marks
  If no working is shown then incorrect (even though nearly correct) answers score no marks.

www.dynamicpapers.com

- **With working**
  If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks.

  If a candidate misreads a number from the question: eg. uses 252 instead of 255; follow through their working and deduct 2A marks from any gained provided the work has not been simplified. (Do not deduct any M marks gained.)

  If there is a choice of methods shown, then award the lowest mark, unless the subsequent working makes clear the method that has been used

  Examiners should send any instance of a suspected misread to review (but see above for simple misreads).

- **Ignoring subsequent work**
  It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. incorrect cancelling of a fraction that would otherwise be correct.

  It is not appropriate ignore subsequent work when the additional work essentially makes t wer incorrect eg algebra.

- **Parts of questions**
  Unless allowed by the mark e, the marks allocated to one part of the question CANNOT be awarded other.

www.dynamicpapers.com

General Principles for Further Pure Mathematics Marking

(but note that specific mark schemes may sometimes override these general principles)

Method mark for solving a 3 term quadratic equation:

1. Factorisation:

$(x^2 + bx + c) = (x + p)(x + q)$, where $|pq| = |c|$ leading to $x = ...$

$(ax^2 + bx + c) = (mx + p)(nx + q)$ where $|pq| = |c|$ and $|mn| = |a|$ leading to $x = ...$

2. Formula:

Attempt to use the **correct** formula (shown explicitly or implied by working) with values for *a*, *b* and *c*, leading to $x = ...$

3. Completing the square:

$x^2 + bx + c = 0$: $\left(x \pm \frac{b}{} \right)^2 \pm q \pm c = 0, \quad q \neq 0$ leading to $x = ...$

Method marks for differe on and integration:

1. Differentiation

Power of at least one term de d by 1. $\left(x^n \to x^{n-1}\right)$

2. Integration:

Power of at least one term increased $\left(x^n \to x^{n+1}\right)$

Use of a formula:

Generally, the method mark is gained by **either**

quoting a correct formula and attempting to use it, even if there are mistakes in the substitution of values

or, where the formula is not quoted, the method mark can be gained by implication from the substitution of correct values and then proceeding to a solution.

www.dynamicpapers.com

Answers without working:

The rubric states "Without sufficient working, correct answers may be awarded no marks".

General policy is that if it could be done "in your head" detailed working would not be required. (Mark schemes may override this eg in a case of "prove or show...."

Exact answers:

When a question demands an exact answer, all the working must also be exact. Once a candidate loses exactness by resorting to decimals the exactness cannot be regained.

Rounding answers (where accuracy is specified in the question)

Penalise only once per ion for failing to round as instructed - ie giving more digits in the answers. Ans ith fewer digits are automatically incorrect, but the is rule may allow the ma e awarded before the final answer is given. before the final answer is given.

www.dynamicpapers.com

MARK SCHEME

Question number	Scheme	Marks
1 (a)	$l = r\theta \Rightarrow l = 13 \times 2 = 26$ (cm)	B1 [1]
(b)	$A = \frac{\theta}{2}r^2 = \frac{2}{2} \times 13^2 = 169$ (cm^2)	M1A1 [2]
		Total 3 marks
(a) **B1**	$l = 26$	
(b) **M1**	Use of $A = \frac{\theta}{2}r^2$ or $A = \frac{1}{2}rl$ or $A = \frac{l^2}{2\theta}$	
A1	$A = 169$	

Question number	Scheme	Marks
2 (a)	Line l_1 $m = \frac{-8}{4} = -2$ $-8 = -2(x) \Rightarrow y + 2x = 8$	M1A1
	Line l_2 $m = \frac{-4}{6} = -\frac{2}{3}$ 4 2 $x) \Rightarrow 3y + 2x = 12$	A1
		[3]
(b)	$x \geq 0$ $y + 2x \leq 8$ $3y + 2x \geq 12$ Accept < a >	B1B1ftB1ft [3]
		Total 6 marks
(a)		
M1	Calculating the gradient of either l_1 or l_2	
A1	$y + 2x = 8$	
A1	$3y + 2x = 12$	
	NB If both are correct but not in the form $ax + by = c$ then award A1A0	
(b)	For all 3 marks accept < and > instead of $\leq$ and $\geq$	
B1	$x \geq 0$	
B1ft	$y + 2x \leq 8$ oe (ft l_1)	
B1ft	$3y + 2x \geq 12$ oe (ft l_2)	

www.dynamicpapers.com

Question number	Scheme	Marks
3	$33 = \frac{1}{2} \times 11 \times 12 \times \sin \angle ABC \Rightarrow \sin \angle ABC = \frac{1}{2}$ $\left[30^\circ \text{ or } 150^\circ\right]$	M1A1
	$AC = \sqrt{11^2 + 12^2 - 2 \times 11 \times 12 \cos 30^\circ} = 6.0306... \approx 6.03$ (cm) $AC = \sqrt{11^2 + 12^2 - 2 \times 11 \times 12 \cos 150^\circ} = 22.2178... \approx 22.2$ (cm)	M1A1 A1 [5]
		Total 5 marks

M1	Use of $\frac{1}{2}ab\sin C = 33$ (must be set = 33)
A1	30° and 150° (Accept $\frac{\pi}{6}$ and $\frac{5\pi}{6}$)
M1	Use of $c^2 = {}^2 \quad b \quad - 2ab\cos C$ (square root not required for M1)
A1	6.03
A1	22.2
	NB If both answers given to 3sf but are correct then award A1A0

www.dynamicpapers.com

Question number	Scheme	Marks
4 (a)	$A = \frac{1}{2} \times 8 \times 8 \times \sin 60^\circ = \left(16\sqrt{3}\right)$	M1
	$48\sqrt{3} = \frac{1}{3} \times '16\sqrt{3}' \times h \Rightarrow h = 9$ *	M1 A1cso [3]
(b)	$BX = \sqrt{9^2 + 8^2} = \sqrt{145}$	M1
	$\angle BXC = \frac{'145' + '145' - 8^2}{2 \times '\sqrt{145}' \times '\sqrt{145}'} = 38.8025...^\circ \approx 38.8^\circ$	M1A1 [3]
(c)	Let midpoint of BC be M	
	$AM = \sqrt{8^2 - 4^2} = \left(4\sqrt{3}\right)$ or $MX = \sqrt{145 - 4^2} = \left(\sqrt{129}\right)$	M1
	$\angle XMA = \tan^{-1}\left(\frac{9}{'4\sqrt{3}'}\right)$ or $\angle XMA = \sin^{-1}\left(\frac{9}{'\sqrt{129}'}\right)$ or $\angle XMA = \cos^{-1}\left(\frac{'4\sqrt{3}'}{'\sqrt{129}'}\right)$	M1
	$= 52.4109...^\circ \quad 52.4$	A1 [3]
	ALT 1	
	Let midpoint of BC be M	
	$(XA)^2 = (AM)^2 + (XM) \quad 2(\quad M)\cos\theta$	{M1}
	$9^2 = \left(8^2 - 4^2\right) + \left(8^2 + 9^2 - 4^2\right) - \sqrt{\quad}\ ^2\sqrt{8^2 + 9^2 - 4^2}\cos\theta$	{M1}
	$\theta = 52.4109...^\circ \approx 52.4^\circ$	{A1} [3]
		Total 9 marks

(a)	
M1	Use of $\frac{1}{2}ab\sin C$ (may be implied by $\left(16\sqrt{3}\right)$
M1	Use of $\frac{1}{3} \times$ 'Area of base' $\times h$
A1 cso	Obtains the **given** answer with no errors in the working
(b)	
M1	Use of $\sqrt{(\text{part } a)^2 + 8^2}$ (may be implied by $\sqrt{145}$)
M1	Use the cosine rule, either form. If not for angle BXC there must be a complete method shown for obtaining BXC (follow through their BX)
A1	awrt 38.8°
(c)	
M1	Use of Pythagoras' to find the length of AM or MX (may be implied by $4\sqrt{3}$ or $\sqrt{129}$)
M1	$\tan^{-1}\left(\frac{9}{'4\sqrt{3}'}\right)$ or $\sin^{-1}\left(\frac{9}{'\sqrt{129}'}\right)$ or $\cos^{-1}\left(\frac{'4\sqrt{3}'}{'\sqrt{129}'}\right)$
A1	awrt 52.4°

www.dynamicpapers.com

ALT	
M1	Use of cosine rule using *AX*, *AM* and *XM* e.g. $(XA)^2 = (AM)^2 + (XM)^2 - 2(AM)(XM)\cos\theta$
M1	Correct values substituted into the cosine rule in any form e.g. $9^2 = \left(8^2 - 4^2\right) + \left(8^2 + 9^2 - 4^2\right) - 2\sqrt{8^2 - 4^2}\sqrt{8^2 + 9^2 - 4^2}\cos\theta$
A1	awrt 52.4°

www.dynamicpapers.com

Question number	Scheme	Marks
5 (a)	$(\alpha+\beta)^3=\alpha^3+3\alpha^2\beta+3\alpha\beta^2+\beta^3 \Rightarrow \alpha^3+\beta^3$ $=(\alpha+\beta)^3-3\alpha\beta(\alpha+\beta)$*	M1A1 cso [2]
(b)	$\alpha+\beta=-\frac{3}{2} \quad \alpha\beta=\frac{6}{2}=3$ $\left[\alpha^3+\beta^3=(\alpha+\beta)^3-3\alpha\beta(\alpha+\beta)\right]$ $\alpha^3+\beta^3=\left(-\frac{3}{2}\right)^3-3\times3\times\left(-\frac{3}{2}\right)=\frac{81}{8}$	B1 B1 [2]
(c)	$(\alpha^2+\beta^2)^2=\alpha^4+2\alpha^2\beta^2+\beta^4 \Rightarrow \alpha^4+\beta^4=(\alpha^2+\beta^2)^2-2(\alpha\beta)^2$*	M1A1 cso [2]
(d)	$\alpha^2+\beta^2=\left(-\frac{3}{}\right)^2 \quad 2\times3=-\frac{15}{4}$ Sum $(\alpha\ -\beta\ ^3\ \alpha)=\alpha^3+\beta^3-(\alpha+\beta)=\frac{81}{8}-\left(-\frac{3}{2}\right)=\frac{93}{8}$ Product $(\alpha^3-\beta)\times(\beta\ \ \beta)^3-(\alpha^4+\beta^4)+\alpha\beta$ $=2\ \left[\left(-\frac{15}{4}\right)^2-2\times3^2\right]+3=\frac{543}{16}$ Equation $x^2-\frac{93}{8}x+\frac{543}{16}=0\Rightarrow16x^2\ \ 18\ \ \ 3=0$	B1 B1 M1 A1 M1A1 [6]
		Total 12 marks

(a)	
M1	$(\alpha+\beta)^3=\alpha^3+3\alpha^2\beta+3\alpha\beta^2+\beta^3$
A1 cso	Obtains the **given** answer with no errors in the working
(b)	
B1	$\alpha+\beta=-\frac{3}{2}$ and $\alpha\beta=\frac{6}{2}=3$
B1	$\alpha^3+\beta^3=\frac{81}{8}$
(c)	
M1	$(\alpha^2+\beta^2)^2=\alpha^4+2\alpha^2\beta^2+\beta^4$
A1 cso	Obtains the **given** answer with no errors in the working

www.dynamicpapers.com

(d)	
B1	$\alpha^2+\beta^2=-\frac{15}{4}\quad\left(\text{May be implied by } \alpha^4+\beta^4=-\frac{63}{16}\right)$
B1	$\left(\alpha^3-\beta\right)+\left(\beta^3-\alpha\right)=\frac{93}{8}$
M1	$\left(\alpha^3-\beta\right)\times\left(\beta^3-\alpha\right)=(\alpha\beta)^3-\left(\alpha^4+\beta^4\right)+\alpha\beta$
A1	$\left(\alpha^3-\beta\right)\times\left(\beta^3-\alpha\right)=\frac{543}{16}$
M1	$x^2-\text{'sum'}x+\text{'product'}(=0)$
A1	$16x^2-186x+543=0$

www.dynamicpapers.com

Question number	Scheme	Marks
6 (a)	$\left[V=\frac{1}{3}\pi r^2 h\right]$ $\tan 30^\circ=\frac{r}{h}\Rightarrow r=\frac{h}{\sqrt{3}}$ $V=\frac{1}{3}\pi\left(\frac{h}{\sqrt{3}}\right)^2 h\Rightarrow V=\frac{1}{9}\pi h^3$ *	M1 A1 cso [2]
(b)	$\frac{\mathrm{d}V}{\mathrm{d}h}=\frac{\pi h^2}{3}$ $\frac{\mathrm{d}h}{\mathrm{d}t}=\frac{\mathrm{d}h}{\mathrm{d}V}\times\frac{\mathrm{d}V}{\mathrm{d}t}\Rightarrow\frac{\mathrm{d}h}{\mathrm{d}t}=\frac{3}{\pi h^2}\times -0.9=\frac{3}{\pi\times 1.2^2}\times -0.9=-0.59683...$ ≈ -0.597 cm/s [Accept an answ of ± 0.597 cm/s]	M1 M1M1 A1 [4]
		Total 6 marks

(a)	
M1	Finding $r=\frac{h}{\sqrt{3}}$ and s ting into $V=\frac{1}{3}\pi r^2 h$ (Allow $r=h\tan 30^\circ$)
A1 cso	Obtains the **given** answer errors in the working
(b)	
M1	$\frac{\mathrm{d}V}{\mathrm{d}h}=\frac{\pi h^2}{3}$
M1	$\frac{\mathrm{d}h}{\mathrm{d}t}=\frac{\mathrm{d}h}{\mathrm{d}V}\times\frac{\mathrm{d}V}{\mathrm{d}t}$
M1	$\frac{\mathrm{d}h}{\mathrm{d}t}=\frac{3}{\pi h^2}\times\pm 0.9$
A1	$\pm 0.597\ (\mathrm{cm/s})$

www.dynamicpapers.com

Question number	Scheme	Marks
7 (a)	$\frac{ar^6}{ar^3} = r^3 = \frac{\mathrm{e}^{\frac{2x+1}{2}}}{\mathrm{e}^{x+2}} = \frac{\mathrm{e}^x \times \mathrm{e}^{\frac{1}{2}}}{\mathrm{e}^x \times \mathrm{e}^2} = \mathrm{e}^{-\frac{3}{2}} \Rightarrow r = \mathrm{e}^{-\frac{1}{2}}$ *	M1M1A1 cso [3]
	ALT $\frac{e^{x+2}}{r^3} r^6 = \mathrm{e}^{\frac{2x+1}{2}} \Rightarrow r^3 = e^{x+\frac{1}{2}-x+2} \Rightarrow r^3 = e^{-\frac{3}{2}} \Rightarrow r = \mathrm{e}^{-\frac{1}{2}}$ *	{M1} {M1}{A1} cso [3]
(b)	$ar^3 = a\mathrm{e}^{-\frac{3}{2}} = \mathrm{e}^{x+2} \Rightarrow a = \frac{\mathrm{e}^{x+2}}{\mathrm{e}^{-\frac{3}{2}}} = \frac{\mathrm{e}^x \times \mathrm{e}^2}{\mathrm{e}^{-\frac{3}{2}}} = \mathrm{e}^{x+\frac{7}{2}}$ oe	M1M1A1 [3]
	ALT $a = \frac{e^{x+2}}{e^{-\frac{3}{2}}} = e^{x+2+\frac{3}{2}} = e^{x+\frac{7}{2}}$ oe	{M1}{M1} {A1} [3]

(a)	
M1	Use of $\frac{ar^6}{ar^3} = r^3$
M1	Using $e^{a+b} = e^a \times e^b$ to simp ding to $r^3 = \mathrm{e}^c$ where c is a number
A1 cso	Obtains the **given** answer with rs in the working
ALT	
M1	For rearranging to make a subject an ituting into the other equation
M1	Using $e^{a-b} = e^a \div e^b$ to simplify leading e^c where c is a number
A1 cso	Obtains the **given** answer with no errors in rking
(b)	
M1	For $a = \frac{e^{x+2}}{e^{-\frac{3}{2}}}$
M1	Using $e^{a+b} = e^a \times e^b$ to simplify to $a = \frac{\mathrm{e}^x \times \mathrm{e}^2}{\mathrm{e}^{-\frac{3}{2}}}$
A1	$\mathrm{e}^{x+\frac{7}{2}}$ oe
ALT	
M1	For $a = \frac{e^{x+2}}{e^{-\frac{3}{2}}}$
M1	Using $e^{a+b} = e^a \times e^b$ to simplify to $e^{x+2+\frac{3}{2}}$
A1	$\mathrm{e}^{x+\frac{7}{2}}$ oe

www.dynamicpapers.com

(c)	$S_\infty = \dfrac{e^{x+\frac{7}{2}}}{1-e^{-\frac{1}{2}}} = \dfrac{e^{x+\frac{7}{2}}}{\frac{e^{\frac{1}{2}}-1}{e^{\frac{1}{2}}}} = \dfrac{e^{x+4}}{e^{\frac{1}{2}}-1} \Rightarrow p = x+4$	M1M1A1 [3]
	ALT 1	
	$S_\infty = \dfrac{e^{x+\frac{7}{2}}}{1-e^{-\frac{1}{2}}} = \dfrac{e^{x+\frac{7}{2}}}{1-e^{-\frac{1}{2}}} \times \dfrac{e^{\frac{1}{2}}}{e^{\frac{1}{2}}} = \dfrac{e^{x+4}}{e^{\frac{1}{2}}-1} \Rightarrow p = x+4$	{M1}{M1} {A1} [3]
	ALT 2	
	$S_\infty = \dfrac{e^{x+\frac{7}{2}}}{1-e^{-\frac{1}{2}}} = \dfrac{e^{p}}{e^{\frac{1}{2}}-1} \Rightarrow e^{x+\frac{7}{2}}\left(e^{\frac{1}{2}}-1\right) = e^{p}\left(1-e^{-\frac{1}{2}}\right)$	{M1}{M1} {A1} [3]
	$\Rightarrow e^{x+4} - e^{x+\frac{7}{2}} = e^{p} - e^{p-\frac{1}{2}} \Rightarrow p = x+4$	

(c)	
M1	Use of $S_\infty = \dfrac{a}{1-r}$
M1	$1-e^{-\frac{1}{2}} = \dfrac{e^{\frac{1}{2}}-1}{e^{\frac{1}{2}}}$
A1	$p = x+4$
ALT 1	
M1	Use of $S_\infty = \dfrac{a}{1-r}$
M1	Multiplying S_∞ by $\dfrac{e^{\frac{1}{2}}}{e^{\frac{1}{2}}}$
A1	$p = x+4$
ALT 2	
M1	Use of $S_\infty = \dfrac{a}{1-r}$
M1	For simplifying to $e^{x+4} - e^{x+\frac{7}{2}} = e^{p} - e^{p-\frac{1}{2}}$ oe
A1	$p = x+4$

www.dynamicpapers.com

(d)	$e^{x+\frac{7}{2}} \times \left(e^{-\frac{1}{2}}\right)^{17} > 1.6 \Rightarrow e^x \times e^{\left(\frac{7}{2}-\frac{17}{2}\right)} > 1.6 \Rightarrow e^x > \frac{1.6}{e^{-5}}$	M1
	$\Rightarrow e^x > 237.46105....$	M1
	$\Rightarrow x > \ln(237.46105) \Rightarrow x > 5.4700...$	M1
	$\Rightarrow x = 6$	A1
		[4]
	ALT	
	$e^{x+\frac{7}{2}} \times \left(e^{-\frac{1}{2}}\right)^{17} > 1.6 \Rightarrow e^{x-5} > 1.6$	{M1}
	$\Rightarrow x - 5 > \ln(1.6)$	{M1}
	$\Rightarrow x > \ln(1.6) + 5 \Rightarrow x > 5.4700...$	{M1}
	$\Rightarrow x = 6$	{A1}
		[4]
		Total 13 marks

(d)	**NB** Allow use of = f arks
M1	Use of $ar^{17} > 1.6$
M1	Simplifying to $e^x > \ldots$
M1	Using logs to reach $x > \ln \ldots$
A1	$x = 6$
ALT	
M1	Use of $ar^{17} > 1.6$
M1	Simplifying to $e^{x-5} > \ldots$ ft their a
M1	Using logs to reach $x > \ln \ldots + 5$ ft their a
A1	$x = 6$

www.dynamicpapers.com

Question number	Scheme	Marks
8 (a)	$k = 2$	B1 [1]
(b)	$2 + 3\cos A - \sin A - 3\sin 2A - 2\cos^2 A = 3\cos A - \sin A - 6\sin A\cos A + 2\sin^2 A$ $3\cos A - \sin A - 6\sin A\cos A + 2\sin^2 A = 3\cos A(1 - 2\sin A) - \sin A(1 - 2\sin A)$ $3\cos A(1 - 2\sin A) - \sin A(1 - 2\sin A) = (1 - 2\sin A)(3\cos A - \sin A)$ $\Rightarrow p = 3,\ q = 1,\ r = 2$	B1 M1 A1 [3]
	ALT	
	$2 + 3\cos A - \sin A - 3\sin 2A - 2\cos^2 A = 3\cos A - \sin A - 6\sin A\cos A + 2\sin^2 A$ $(p\cos A - \sin A)(q - r\sin A) = pq\cos A - q\sin A - pr\cos A\sin A + r\sin^2 A$ $pq\cos A - q\sin A - pr\cos A\sin A + r\sin^2 A$ $\Rightarrow p = 3,\ q = 1,$	{B1} {M1} {A1} [3]
(c)	$(1 - 2\sin 2\theta)(3\cos 2\quad\theta) = 0$ $\Rightarrow 3\cos 2\theta - \sin 2\theta\quad 0\qquad = 3$ $\Rightarrow 1 - 2\sin 2\theta = 0 \Rightarrow \sin 2\quad = \frac{}{2}$ $\tan 2\theta = 3 \Rightarrow 2\theta = 1.2490...,\ 4.3906.\qquad \theta = 0.625,\ 2.20$ $\sin 2\theta = \frac{1}{2} \Rightarrow 2\theta = 0.5235...,\ 2.6179...\qquad 62,\ 1.31\ \left[\frac{\pi}{12}, \frac{5\pi}{12}\right]$	B1 B1 M1A1 M1A1 [6]
		Total 10 marks

(a)

B1 $k = 2$

(b)

B1 Substituting $k = 2$ and use of $\sin^2 A + \cos^2 A = 1$ to obtain $3\cos A - \sin A - 6\sin A\cos A + 2\sin^2 A$

M1 Factorising to obtain $(1 - 2\sin A)(3\cos A - \sin A)$

A1 $p = 3,\ q = 1,\ r = 2$ If p, q and r are stated then they must be correct (may be implied by a correct factorisation if p, q and r are not stated)

ALT

B1 Substituting $k = 2$ and use of $\sin^2 A + \cos^2 A = 1$ to obtain $3\cos A - \sin A - 6\sin A\cos A + 2\sin^2 A$

M1 Expanding $(p\cos A - \sin A)(q - r\sin A)$ to obtain $pq\cos A - q\sin A - pr\cos A\sin A + r\sin^2 A$

A1 $p = 3,\ q = 1,\ r = 2$

www.dynamicpapers.com

(c)	
B1	$\tan 2\theta = 3$
B1	$\sin 2\theta = \frac{1}{2}$
M1	$2\theta = 1.24(90...,)\ 4.39(06...\ )$
A1	$\theta = 0.625,\ 2.2(0)$
M1	$2\theta = 0.5235...,\ 2.61(79...)$ allow $\left[\frac{\pi}{6}, \frac{5\pi}{6}\right]$
A1	$\theta = 0.262,\ 1.31$ allow $\left[\frac{\pi}{12}, \frac{5\pi}{12}\right]$

www.dynamicpapers.com

Question number	Scheme	Marks
9 (a)	$\frac{1}{(2-x)^3} = (2-x)^{-3} = \frac{1}{8}\left(1-\frac{x}{2}\right)^{-3} \Rightarrow p = \frac{1}{8},\ q = \frac{1}{2}$	B1B1 [2]
(b)	$\frac{1}{8}\left(1-\frac{x}{2}\right)^{-3} = \frac{1}{8}\left[1+(-3)\left(-\frac{x}{2}\right)+\frac{(-3)(-4)\left(-\frac{x}{2}\right)^2}{2!}+\frac{(-3)(-4)(-5)\left(-\frac{x}{2}\right)^3}{3!}\right]$ $= \frac{1}{8}+\frac{3}{16}x+\frac{3}{16}x^2+\frac{5}{32}x^3+\ldots$	M1 A1A1 [3]
(c)	$(a+bx)\left(\frac{1}{8}+\frac{3}{16}+\frac{3}{6}x^2\right) = \frac{a}{8}+x\left(\frac{2b}{16}+\frac{3a}{16}\right)+\left\{x^2\left(\frac{3a}{16}+\frac{3b}{16}\right)\right\}$ $\Rightarrow \frac{3}{8} = \frac{a}{8} \Rightarrow a = 3$ $\Rightarrow -\frac{43}{16} = \frac{2b+3a}{16} \Rightarrow 2\ =-\quad -52 \Rightarrow b = -26$	M1 A1 A1 [3]
(d)	$\frac{3a+3b}{16} = \frac{9-78}{16} = -\frac{69}{16}$ oe	M1A1 [2]
		Total 10 marks

	NB If p and q are stated then they must be correct t if p and q are not stated then
(a)	$\frac{1}{8}\left(1-\frac{x}{2}\right)^{-3}$ scores B1B1
B1	$p = \frac{1}{8}$ (Allow $p = 2^{-3}$)
B1	$q = \frac{1}{2}$

www.dynamicpapers.com

(b)	
M1	Attempts to use the binomial expansion for their $(1-qx)^{-3}$. Must have first term 1, three more terms with ascending powers of x, 2 or 2! and 6 or 3! seen , and their $\left(-\frac{x}{2}\right)$ used at least once. No simplification needed. Ignore terms beyond x^3
A1	Two algebraic terms correct in the expansion for their $(1-qx)^{-3}$ .. Must be single fractions, not necessarily in lowest terms. Ignore terms beyond x^3
A1	All four terms correct and in lowest terms. Ignore terms beyond x^3
(c)	
M1	For either their $\frac{1}{8}a=\frac{3}{8}$ or their $\frac{3}{16}a+$ their $\frac{1}{8}b=-\frac{43}{16}$ May be implied by a correct value of a or b.
A1	$a=3$
A1	$b=-26$ **NB** answers a nd $b=-26$ scores 3/3
(d)	
M1	Substituting their a ir b into their $\frac{3a+3b}{16}$
A1	$-\frac{69}{16}$ oe
	NB If $p=2,\ q=\frac{1}{2}$ is used then and $b=-\frac{13}{8}$ which substituted into $3a+3b$ gives an answer of $-\frac{69}{16}$ but scores A0

www.dynamicpapers.com

Question number	Scheme	Marks
10 (a)	$\int\left(3x^2-4x-p\right)\,\mathrm{d}x=\frac{3x^3}{3}-\frac{4x^2}{2}-px+c\quad\left[=x^3-2x^2-px+c\right]$	M1A1
	$y=x^3-2x^2-px+c$	
	At $(2,0)\quad 0=8-8-2p+c\Rightarrow c=2p$	M1
	At $(-1,9)\quad 9=-1-2+p+c\Rightarrow c=12-p$	M1
	$\Rightarrow p=4, c=8\Rightarrow y=x^3-2x^2-4x+8$*	A1A1cso [6]
(b)	$x^3-2x^2-4x+8=8-4x\Rightarrow x^3-2x^2=0\Rightarrow x^2(x-2)=0$	M1
	$x=0, x=2$	A1
	Area $=$ ^{2} 8 $)\,\mathrm{d}x-\int_0^2\left(x^3-2x^2-4x+8\right)\mathrm{d}x$	M1
	Area $=$ $(8\quad 4x\qquad \int_0^2\left(x^3-2x^2-4x+8\right)\mathrm{d}x=\int_0^2\left(-x^3+2x^2\right)\mathrm{d}x$	
	Area $=\left[-\frac{x^4}{4}+\frac{x}{3}\right.^{2}_{0}\qquad\left.\frac{16}{}\right)-(0)=\frac{4}{3}$	M1M1A1 [6]
		Total 12 marks

(a)

M1 Attempts to integrate

A1 Correct integration including + *c*

M1 Substitution of (2, 0) (Does not have to be s ed)

M1 Substitution of (–1, 9 (Does not have to be sim)

A1 $p=4, c=8$

A1 cso Obtains the **given** answer with no errors in the working

(b)

M1 Equating *C* and *l*

A1 $x=0$ and $x=2$

NB If correct limits are seen then M1A1 is awarded

M1 Use of $\int_a^b\left(\mathrm{f}(x)-\mathrm{g}(x)\right)\mathrm{d}x$ or $\int_a^b\mathrm{f}(x)\mathrm{d}x-\int_a^b\mathrm{g}(x)\mathrm{d}x$ or $\int_a^b\mathrm{f}(x)\mathrm{d}x-\frac{1}{2}\times 2\times 8$

Ignore limits (These can be either way round)

M1 Attempt the integration. Limits not needed.

M1 Substitute the correct limits. $\left(\text{May be implied by } \pm\frac{4}{3}\right)$

A1 $\frac{4}{3}$

NB If no integration is seen then M0M0A0 is awarded for the last 3 marks for an answer of $\frac{4}{3}$

www.dynamicpapers.com

Question number	Scheme	Marks
11 (a)	(i) $y=3$ (ii) $x=-1$	B1B1 [2]
(b) (i) (ii)	$0=\frac{3x-2}{x+1}\Rightarrow 3x=2\Rightarrow x=\frac{2}{3}$ $y=\frac{3\times 0-2}{0+1}=-2$	B1 B1 [2]
(c)	 http://britishstudent.com.wordpress.com/	B1 (shape and position) B1 (asymptotes) B1 (intersections) [3]
(d)	$mx+4=\frac{3x-2}{x+1}\Rightarrow mx^2+x(m+1)+6=0$ $b^2-4ac<0\Rightarrow (m+1)^2-4\times m\times 6<0\Rightarrow m^2-22m+1<0$ $(m-11)^2-120=0\Rightarrow m=11\pm\sqrt{120}$ Defines region $11-2\sqrt{30}<m<11+2\sqrt{30}$	M1 M1 M1A1 M1 depM1A1 [7]
		Total 14 marks

www.dynamicpapers.com

(a) (i)	
B1	$y = 3$
(a) (ii)	
B1	$x = -1$
(b) (i)	
B1	$x = \frac{2}{3}$
(b) (ii)	
B1	$y = -2$
(c)	
B1	Two curves in the correct quadrant (Condone poor shape as long as intention is clear)
B1	Two asymptotes drawn and correctly labelled (Allow –1 and 3 correctly labelled on axes to count as labelled). There must be at least one part of the curve drawn
B1	Intersections labelled
(d)	
M1	Equates *C* and ndone use of ≠ and <)
M1	Forms a 3TQ (C the use of ≠ and <) (allow 1 error in the expansion of correct brackets or in simplifying)
M1	Use of $b^2 - 4ac <$ to 3TQ
A1	Simplify to $m^2 - 22$ 1
M1	Solving the 3TQ If the quad incorrect then the method for solving must be shown
depM1	Simplifying to the form $a \pm b\sqrt{2}$ D ant on previous M1
A1	$11 - 2\sqrt{30} < m < 11 + 2\sqrt{30}$ (Allow a = 30)

www.dynamicpapers.com

www.dynamicpapers.com

www.dynamicpapers.com

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom