

Mark Scheme (Results)

January 2016

Pearson Edexcel International A Level in Statistics 2 (WST02)
Paper 01

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at <a href="https://www.edexcel.com">www.edexcel.com</a> or <a href="https://www.edexcel.com">www.btec.co.uk</a>. Alternatively, you can get in touch with us using the details on our contact us page at <a href="https://www.edexcel.com/contactus">www.edexcel.com/contactus</a>.

#### Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at:

www.pearson.com/uk

January 2016
Publications Code IA043349
All the material in this publication is copyright
© Pearson Education Ltd 2016

## **General Marking Guidance**

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

#### PEARSON EDEXCEL IAL MATHEMATICS

### **General Instructions for Marking**

- 1. The total number of marks for the paper is 75
- 2. The Edexcel Mathematics mark schemes use the following types of marks:
- M marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
- A marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
- **B** marks are unconditional accuracy marks (independent of M marks)
- Marks should not be subdivided.

#### 3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod benefit of doubt
- ft follow through
- the symbol  $\sqrt{}$  will be used for correct ft
- cao correct answer only
- cso correct solution only. There must be no errors in this part of the question to obtain this mark
- isw ignore subsequent working
- awrt answers which round to
- SC: special case
- oe or equivalent (and appropriate)
- d... or dep dependent
- indep independent
- dp decimal places
- sf significant figures
- \* The answer is printed on the paper or ag- answer given
- or d... The second mark is dependent on gaining the first mark
- 4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.
- 5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.

- 6. If a candidate makes more than one attempt at any question:
  - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
  - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
- 7. Ignore wrong working or incorrect statements following a correct answer.

# **Special notes for marking Statistics exams**

- If a candidate is "hedging their bets" e.g. give Attempt 1...Attempt 2...etc then please send to review.
- Any correct method should gain credit. If you cannot see how to apply the mark scheme but believe the method to be correct please send to review.

Question Number	Scheme	Marks
1 (a)	Any suitable problem with using the database as a sampling frame:  E.g.  Might not be up to date  Might not contain all customers (incomplete)  Might be biased	B1
		(1)
<b>(b)</b>	B(20, 0.35)	B1 dB1
		(2)
(c)	$P(X > 10) = 1 - P(X \le 10) = 1 - 0.9468 = 0.0532$ awrt <b>0.0532</b>	M1 A1 (2) Total (5)
	Notes	
(a)	B1 Ignore extraneous non-contradictory comments 'Not accurate' without context is not sufficient	
<b>(b)</b>	$1^{\text{st}}$ B1 for B or binomial which must be seen in part (b) $2^{\text{nd}}$ B1 dependent on $1^{\text{st}}$ B1 for ([ $n = ]20$ , [ $p = ]0.350e$ ) which must be seen in part (b)	
(c)	M1 for writing or using $1-P(X \le 10)$	

Question Number	Scheme	Marks	
2 (a)	$\frac{5-3}{b-a} = \frac{1}{8} \Rightarrow b-a = 16$ $\frac{a+b}{2} = 4 \Rightarrow a+b = 8$	M1 M1	
	a = -4 and $b = 12$	A1 (3)	
(b)	$[cE(X) - 2 =] 4c - 2 = 0$ or $\int_{-4}^{12} \frac{1}{16!} (cx - 2) dx = 0$	M1	
	$c = \frac{1}{2}$	A1 (2)	
(c)	$Var(X) = \frac{(12' - (-4'))^2}{12} \left( = \frac{64}{3} \right) \qquad \underline{or} \qquad \int_{-4'}^{12'} \frac{1}{16'} x^2 dx$	B1ft	
	$E(X^{2}) = \left(\frac{64}{3}\right) + 4^{2} = \frac{112}{3}$ or $\left[\frac{12^{13}}{48} - \frac{-4^{13}}{48}\right] = \frac{112}{3}$	M1 A1	
(d)	P(2X > a+b) = P(X > 4) = 0.5	(3) M1 A1 (2) Total (10)	
	Notes		
(a)			
	A1 both $a = -4$ and $b = 12$		
(b)	M1 for $4c - 2 = 0$ or attempt correct equation for $E(cX - 2) = 0$ using integration		
(c)	<ul> <li>B1ft for substituting their a and b into a correct expression for Var(X) <ul> <li>or correct integral with limits (ft their a and b) for E(X²)</li> </ul> </li> <li>M1 for substituting E(X) = 4 and their Var(X) into a correct expression for E(X²) <ul> <li>or correct integration with correct use of limits (ft their a and b)</li> </ul> </li> <li>A1 for 112/3 or exact equivalent</li> </ul>		
(d)	M1 for correct rearrangement up to $P(X > k)$ where $k = \frac{a+b}{2}$ (may be implied by a correct ft answer)		

	uestion umber	Scheme	Marks
3	(a)(i)	B(60, 0.1)	B1
		$[P(Y=0) + P(Y=1) = ](0.9)^{60} + 60(0.1)^{1}(0.9)^{59}$	M1
	(ii)	0.013777 awrt <u>0.0138</u>	A1
		<u></u>	(3)
	<b>(b)</b>	Po(6)	B1
	, ,	D/W < 1) 0.01725	B1
		$P(Y \le 1) = 0.01735$ or 0.0174 from tables awrt <b>0.0174</b>	ВІ
			(2)
	(c)	N(6, 5.4)	B1
	(C)		
		$P(Y \le 1) \approx P\left(Z < \frac{1.5 - 6}{\sqrt{5.4}}\right)$	M1 dM1
		$P(Y \le 1) \approx P(Z < \frac{1}{\sqrt{5.4}})$	
		= P(Z < -1.936)	A1
			A1 (5)
		= 1 - 0.9738 = 0.0262 (or from calc 0.02640)	(5)
	( T)		D.1
	<b>(d)</b>	(Poisson approximation is more suitable since)	B1
		E.g.	(1)
		n is large and p is small	
		p is not close to 0.5	
		(b) is closer to the true value	Total (11)
			10tai (11)
Notes			
	(a)	Mark (i) and (ii) together	
1		R1 for writing or using $R(60, 0.1)$	

B1 for writing or using B(60, 0.1)

M1 for correct expression (allow use of 60C0 and 60C1)

(Special Case: if no expression given, then answer only scores B1 M0 A0)

- **(b)** 1<sup>st</sup> B1 for writing or using Po(6) (may be implied by a correct answer) 2<sup>nd</sup> B1 awrt 0.0174
- mean = 6 and variance = 5.4 (may be seen in standardisation) **(c)**

$$1^{\text{st}} \, \text{M1} \, \pm \left( \frac{1.5 \, \text{or} \, 1 \, \text{or} \, 0.5 - \textit{their mean}}{\textit{their sd}} \right)$$
 $2^{\text{nd}} \, \text{M1} \, \text{dependent on} \, 1^{\text{st}} \, \text{M1 for using a continuity correction} \, 1 \pm 0.5$ 

$$1^{\text{st}} \text{ A} = \frac{1.5 - 6}{\sqrt{5.4}} \text{ or awrt } -1.94$$

2<sup>nd</sup> A1 awrt 0.026

B1 for a correct supporting reason (d) (Condone n is large together with np < 10) (Condone mean (6)  $\approx$  variance (5.4))

Question Number	Scheme	Mark	S
4 (a)	F(2) = 1 or verify		
	$\begin{vmatrix} \frac{1}{20}(d^4) + \frac{1}{5} = 1 & \frac{1}{20}(2^4) + \frac{1}{5} = 1 \\ d = 2 & \therefore d = 2 \end{vmatrix}$	M1 A1 cso	(2)
(b)	$F(1.5) = \frac{1}{20}(1.5)^4 + \frac{1}{5} = \frac{29}{64}$ awrt <b>0.453</b>	M1 A1	(2)
(c)	1	B1	
(d)	$\frac{1}{20}x^4 + \frac{1}{5} = 0.5$	M1 M1A1	(1)
	$x^4 = 6 \Rightarrow x = 1.56508$ awrt <u>1.57</u>	1,1111	(3)
(e)	P(X > 1.9) = 1 - F(1.9) = 1 - 0.851605 = 0.148395 P(X < k) = 0.148395	M1 A1	
	$\frac{1}{4}k = 0.148395$	M1	
	k = 0.59358 awrt <b>0.594</b>	A1	(4)
		Total (12	2)
	Notes		
(a)	M1 equating $F(d) = 1$ or substituting $d = 2$ and verifying A1cso for correct conclusion with no incorrect working seen (must reject $d = -2$ if given)		
(b)	M1 for substituting 1.5 into third line of $F(x)$ on its own		
(d)	1 <sup>st</sup> M1 setting third line of $F(x) = 0.5$ 2 <sup>nd</sup> M1 solving equation as far as $x^4 = k$		
(e)	$1^{\text{st}}$ M1 for writing or using $1 - F(1.9)$ $1^{\text{st}}$ A1 awrt 0.148 (may be implied by $k = \text{awrt } 0.59$ )		
	$2^{\text{nd}}$ M1 for writing or using $\frac{1}{4}k = 1 - \text{F}(1.9)$		
	2 <sup>nd</sup> A1 awrt 0.594		

Question Number	Scheme		Marks	5
5 (a)	[Po(1)] $(P(X \ge 1))^2 = (1 - 0.3679)^2 = 0.39955041$	awrt <u><b>0.400</b></u>	M1A1	(2)
(b)	Po(2) P( $X = 0$ ) = 0.1353	awrt <u><b>0.135</b></u>	B1 B1	(2)
(c)	P(X = 4) = P(X \le 4) - P(X \le 3)[= 0.0443] Using tables If $\frac{w}{10} = 8$ , P(X \le 4) - P(X \le 3) = 0.0996 - 0.0424 = 0.0572 If $\frac{w}{10} = 8.5$ , P(X \le 4) - P(X \le 3) = 0.0744 - 0.0301 = 0.0443 $\frac{w}{10} = 8.5$ ,		M1	
	so $w = 85$ .		A1	(3)
(d) (e)	[H <sub>0</sub> :] $\mu = 10$ ( $\lambda = 1$ ) $P(X \le 14   \mu = 10) = 0.9165$ $P(X \le 15   \mu = 10) = 0.9513$ So critical region is $X \ge 16$		M1 A1	(1) (2) (2)
	Notes		10141 (10	
(a)	M1 using $(1 - P(X = 0))^2$			
(b)	1 <sup>st</sup> B1 writing or using Po(2) (may be implied by sight of e <sup>-2</sup> )			
(c)	M1 writing or using of $P(X = 4) = P(X \le 4) - P(X \le 3)$ 1 <sup>st</sup> A1 for 8.5 2 <sup>nd</sup> A1 for 85			
(d)	Must use $\lambda$ or $\mu$ with either = 10 or = 1 (Ignore H <sub>1</sub> if given)			
(e)	M1 for using Po(10) (may be implied by e.g. $P(X \ge 15) = 0.0835$ or $P(X \ge 16) = 0.0487$ ) A1 for $X \ge 16$ or $X > 15$ (allow any letter for $X$ ) $P(X \ge 16)$ is A0 A two-tailed critical region is A0			

Question Number	Scheme	Marks
6 (a)	$\int_{1}^{7} (ax^2 + bx) dx = 1$	M1
	$\left[\frac{ax^3}{3} + \frac{bx^2}{2}\right]^7 [=1]$	A1
	$\left(\frac{a(7^3)}{3} + \frac{b(7^2)}{2}\right) - \left(\frac{a}{3} + \frac{b}{2}\right) = 1$	dM1
	$\begin{pmatrix} 3 & 2 \end{pmatrix} \begin{pmatrix} 3 & 2 \end{pmatrix}$ 114a + 24b = 1*	A1cso
	$11\pi u + 2\pi v - 1$	(4)
(b)	$114(\frac{1}{90}) + 24b = 1 \Longrightarrow b = -\frac{1}{90}$	B1
	$[E(X) = ] \int_{1}^{7} (\frac{1}{90} x^{3} + b' x^{2}) dx$	M1
	$\left[\frac{x^4}{360} - \frac{x^3}{270}\right]^7, = \left(\frac{7^4}{360} - \frac{7^3}{270}\right) - \left(\frac{1}{360} - \frac{1}{270}\right) = 5.4$	A1ft, A1oe
		(4)
(c)	$\int_{1}^{x} \left(\frac{1}{90}t^{2} - \frac{1}{90}t\right) dt = \left[\frac{t^{3}}{270} - \frac{t^{2}}{180}\right]^{x}  \text{or } \int \left(\frac{1}{90}x^{2} - \frac{1}{90}x\right) dx = \frac{x^{3}}{270} - \frac{x^{2}}{180} + c \text{ with } F(1) = 0 \text{ or } F(7) = 1$	M1
	$F(x) = \begin{cases} 0 & x < 1\\ \frac{x^3}{270} - \frac{x^2}{180} + \frac{1}{540} & 1 \le x \le 7\\ 1 & x > 7 \end{cases}$	A1 B1
	(-	(3)
(d)	P(X > 5.4') = 1 - F(5.4') = 1 - 0.42305 = 0.5769 awrt <u><b>0.577</b></u>	M1 A1
(e)	Since (d)>0.5, [the mean is less than the median] therefore negative (skew).	M1, A1 (2)
		Total (15)
	Notes	
(a)	1 <sup>st</sup> M1 attempting to integrate $f(x)$ , (at least one term $x^n \to x^{n+1}$ ). Ignore limits. 1 <sup>st</sup> A1 fully correct integration. Ignore limits and accept any letters. 2 <sup>nd</sup> dM1 dep on 1 <sup>st</sup> M1. Subst in correct limits and equate to 1 2 <sup>nd</sup> A1 correct solution only (answer given)	
(b)	B1 for $b = -\frac{1}{90}$	
	1 <sup>st</sup> M1 attempting to integrate $xf(x)$ , (at least one term $x^n \to x^{n+1}$ ). Ignore limits. 1 <sup>st</sup> A1ft fully correct integration ft their value of $b$ . (Allow if given in terms of $a$ and $b$ ) 2 <sup>nd</sup> A1 5.4 oe	
(c)	M1 attempt to integrate $f(x)$ with correct limits or with + C and attempt at $F(1) = 0$ or $F(7) = 1$	
	A1 $\frac{x^3}{270} - \frac{x^2}{180} + \frac{1}{540}$ with correct limits (allow < or $\leq$ ) B1 for top and bottom lines with correct limits	
(d)	M1 $1 - F(5.4') \text{ or } F(7) - F(5.4') \text{ or } \int_{5.4'}^{7} (\frac{1}{90}x^2 + (-\frac{1}{90})'x) dx$	
(e)	M1 for correctly comparing 'their (d)' with 0.5 (may be implied by a correct comparison of mean and median ft their (d)) If no answer given in (d), then M0.  A1 for negative skew which must follow from 'their (d)' > 0.5	

Question Number	Scheme	Marks
7 (a)	$F \sim \text{Po}(4)$ $P(F \ge 6) = 1 - 0.7851 = 0.2149$ Let $Y = \text{the number of fishing trips where at least 6 fish are caught}$ $Y \sim B(5, 0.2149)$ $P(Y = 3) = 10(0.2149)^3 (0.7851)^2 = 0.06117291856$ awrt <u>0.0612</u>	B1 M1 A1 B1 dM1 A1
(b)	H <sub>0</sub> : $\lambda = 4$ ( $\mu = 8$ ) H <sub>1</sub> : $\lambda \neq 4$ ( $\mu \neq 8$ ) $X \sim \text{Po}(8)$	(6) B1 B1
	$P(X \ge 14) \qquad \text{or CR}$ $=1-P(X \le 13)=1-0.9658$ $=0.0342 \qquad P(X \le 14)=0.9827$ $=0.0342 > 0.025] \text{ so do not reject } H_0 \text{ or } 14 \text{ is not in the critical region } \text{or Not significant}$ There is not enough evidence of a <u>change</u> in the	M1 A1 dM1
	mean number of fish caught or number of fish caught per hour or rate of fish caught.	(6) Total (12)
	Notes	
(a)	1st B1 writing or using Po(4) 1st M1 writing or using $1-P(F \le 5)$ 1st A1 for awrt 0.215 2nd B1 writing or using binomial $(5, p)$ 2nd M1 dependent on $2^{nd}$ B1 correct expression for $P(Y = 3)$ using their value of $p$ 2nd A1 awrt 0.0612	
(b)	1 <sup>st</sup> B1 both hypotheses correct. Must use $\lambda$ or $\mu$ 2 <sup>nd</sup> B1 writing or using Po(8)  1 <sup>st</sup> M1 writing 1−P( $X \le 13$ ) (may be implied by sight of 1 − 0.9658)  or for CR method: P( $X \le 13$ ) = 0.9658 P( $X \le 14$ ) = 0.9827  1 <sup>st</sup> A1 for probability awrt 0.0342 or CR $X > 14$ or $X \ge 15$ 2 <sup>nd</sup> dM1 Dep. on 1 <sup>st</sup> B1. Non-contradictory statement which follows from their probabil 2 <sup>nd</sup> A1cso correct contextual statement and fully correct solution with all other marks so	