

Please check the examination details below before entering your candidate information

Candidate surname					Other names				
Centre Number					Candidate Number				

Pearson Edexcel International Advanced Level

Wednesday 30 October 2024

Morning (Time: 1 hour 20 minutes)

Paper reference **WCH16/01**

Chemistry

International Advanced Level

UNIT 6: Practical Skills in Chemistry II

You must have:
Scientific calculator, ruler

Total Marks

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
– *there may be more space than you need.*

Information

- The total mark for this paper is 50.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*
- You will be assessed on your ability to organise and present information, ideas, descriptions and arguments clearly and logically, including your use of grammar, punctuation and spelling.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P78394A

©2024 Pearson Education Ltd.
V:1/1/1/1/



Answer ALL the questions. Write your answers in the spaces provided.

- 1 (a) Compounds **P** and **Q** are crystalline solids.

P is white and contains one cation and one anion.

Q is pale violet and contains two cations and one anion.

A student carried out tests to identify **P** and **Q**.

- (i) Complete the table to show, by name or formula, the products of the tests.

(4)

Test	Observation	Name or formula of product
Aqueous silver nitrate and dilute nitric acid were added to a sample of an aqueous solution of P Concentrated ammonia solution was then added	A yellow precipitate was formed which was insoluble in concentrated ammonia	The precipitate was
A sample of solid Q was added to 5 cm ³ of aqueous sodium hydroxide and the mixture was warmed	A pungent gas was evolved which turned damp red litmus paper blue A red-brown precipitate was formed which was insoluble in excess alkali	The gas was The precipitate was
Aqueous barium chloride and dilute hydrochloric acid were added to a sample of a solution of Q	A white precipitate was formed	The precipitate was

- (ii) A green flame was seen when a flame test was carried out on a sample of solid **P**. Identify **P** by name or formula.

(1)

P

.....



- (iii) Give the formulae of the three ions present in **Q** and hence the formula of compound **Q**.

(2)

Ions in **Q**

Formula of **Q**

- (b) The student added a sample of a pale-brown, aqueous solution of **Q** to a colourless aqueous solution of **P**. The student observed the formation of a darker brown solution along with a white precipitate.

The student concluded from the formation of the darker brown solution that a redox reaction had taken place.

Explain why the student came to this conclusion.

(2)

(Total for Question 1 = 9 marks)


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



- 2 A group of students investigated the kinetics of the reduction of peroxydisulfate ions by iodide ions in aqueous solution, as shown in Reaction 1.



Some of the iodine produced in Reaction 1 may be removed by including a portion of sodium thiosulfate in the reaction mixture.

The thiosulfate ions react with the iodine as shown in Reaction 2.



After all the thiosulfate ions have reacted, any further iodine produced can be detected by the formation of a starch-iodine complex. Student **M** carried out a sequence of experiments to determine the order of the reaction with respect to **iodide ions**.

The volumes of the solutions used in the mixtures are shown in Table 1.

Table 1

Mixture	Volume KI(aq) / cm ³	Volume K ₂ S ₂ O ₈ (aq) / cm ³	Volume Na ₂ S ₂ O ₃ (aq) / cm ³	Volume H ₂ O(l) / cm ³
1	18.0	20.0	5.0	2.0
2	15.0	20.0	5.0	5.0
3	11.0	20.0	5.0	9.0
4	7.5	20.0	5.0	12.5
5	4.0	20.0	5.0	16.0

The concentrations of the solutions used were

Potassium iodide	KI(aq)	0.200 mol dm ⁻³
Potassium peroxydisulfate	K ₂ S ₂ O ₈ (aq)	0.100 mol dm ⁻³
Sodium thiosulfate	Na ₂ S ₂ O ₃ (aq)	0.00500 mol dm ⁻³

Procedure

Step 1 A few drops of starch solution are added to 5.0 cm³ sodium thiosulfate solution in a small beaker.

Step 2 The potassium peroxydisulfate solution and water are added to the beaker.

Step 3 Then potassium iodide solution is added, the mixture stirred and a timer is started. The time taken to change colour is recorded.

The procedure is repeated for each mixture using the volumes shown in Table 1.



- (a) (i) Suggest a suitable piece of apparatus for measuring the volume of the potassium iodide solution.

(1)

- (ii) State the final colour of the reaction mixture in Step 3.

(1)

- (iii) The reciprocal of the time taken for the mixture to change colour ($1 / \text{time}$), can be used as an approximate measure of the initial rate.

State why the amount of added thiosulfate ions should be much smaller than the amount of peroxydisulfate or iodide ions for this approximation to be valid.

(1)

- (iv) State why is it important that the rate of Reaction 2 is much faster than that of Reaction 1.

(1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(b) The results that Student **M** obtained are shown.

Table 2

Mixture	$[I^-]$ / mol dm ⁻³	time / s	1 / time / s ⁻¹
1	0.080	130	0.0077
2	0.067	157	0.0064
3		202	0.0050
4	0.033	320	0.0031
5	0.018	658	0.0015

- (i) Complete Table 2 with the iodide ion concentration in Mixture 3, using the information from Table 1 and the concentration of the potassium iodide solution.

(1)

- (ii) Plot a graph of 1 / time against the iodide ion concentration.


(2)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



- (iii) Deduce the order of the reaction with respect to iodide ions.

Justify your answer.

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

- (c) (i) Student **N** carried out a sequence of experiments to find the order of reaction with respect to **peroxydisulfate ions**, using solutions of the same concentration.

The volumes of potassium peroxydisulfate solution used were different.

State **two** variables which should be kept the same.

(2)

- (ii) Student **N** found that the reaction is first order with respect to peroxydisulfate ions.

Write the overall rate equation for the reaction of peroxydisulfate ions with iodide ions.

(1)



- (d) Student **R** carried out experiments to find the activation energy of this reaction. A similar procedure was used to determine the rate of reaction at two different temperatures.

The results of these experiments are shown.

$T/^{\circ}\text{C}$	T/K	$1/T/\text{K}^{-1}$	t/s	$k/\text{dm}^3\text{mol}^{-1}\text{s}^{-1}$	$\ln k$
12	285	0.00351	265	2.4×10^{-3}	-6.03
45	318	0.00314	20	3.1×10^{-2}	-3.47

Calculate, without drawing a graph, the activation energy of the reaction.

You should include a sign and units in your answer.

$$\ln k = -\frac{E_a}{R} \times \frac{1}{T} + \text{constant} \quad R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$$


(3)

(Total for Question 2 = 15 marks)



- 3** Ethanedioic acid, $(\text{COOH})_2 \cdot x\text{H}_2\text{O}$, in solution is an acidic bleach for wood.

A student determined the value of x in a sample of ethanedioic acid by titration with potassium manganate(VII) solution.



Procedure

Step 1 A sample of 1.27 g of the solid acid was dissolved in deionised water to make 250.0 cm^3 of solution.

Step 2 25.0 cm^3 of the ethanedioic acid solution was transferred to a conical flask, 25 cm^3 dilute sulfuric acid was added and the mixture warmed to 60°C .

Step 3 The mixture was titrated with potassium manganate(VII) solution of concentration $0.0203\text{ mol dm}^{-3}$.

- (a) (i) Describe how to make 250.0 cm^3 of a solution of ethanedioic acid in Step 1.

(3)

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (ii) Explain why, in Step 2, the ethanedioic acid solution was measured using a 25.0 cm^3 volumetric pipette but the dilute sulfuric acid was measured using a 25 cm^3 measuring cylinder.

(2)

.....

.....

.....

.....

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(iii) State why the titration needed no indicator.

(1)

(iv) Suggest why, in Step 2, the temperature of the mixture at the start should be 60°C but no further warming is needed after a small quantity of the potassium manganate(VII) solution has been added.

(3)

(v) The mean titre was 19.90 cm³.
Determine the value of x in (COOH)₂·xH₂O.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(b) Ethanedioic acid dissociates in two steps.

The **first** dissociation equation is shown.



The acid dissociation constant, $K_{\text{a}1}$, for **this** dissociation is $5.6 \times 10^{-2} \text{ mol dm}^{-3}$.

The corresponding $\text{p}K_{\text{a}1}$ is 1.25.

Outline a practical method that would enable a student to confirm the value for this first acid dissociation constant.

You may use the Indicator data provided.

Assume that equimolar solutions of ethanedioic acid and sodium hydroxide, titration apparatus and a pH meter are available.

You are not expected to include details of any calculations needed.

Indicator data

Indicator	$\text{p}K_{\text{in}}$ (at 298 K)	acid	pH range	alkaline
thymol blue (acid)	1.7	red	1.2–2.8	yellow
methyl orange	3.7	red	3.2–4.4	yellow
bromophenol blue	4.0	yellow	2.8–4.6	blue
phenolphthalein	9.3	colourless	8.2–10.0	red

(3)



(Total for Question 3 = 16 marks)



4 Azo dyes are widely used to dye cotton fabric.

The azo dye, 1-phenylazo-2-naphthol, can be made from phenylamine by reacting it with sodium nitrite and hydrochloric acid.


The diazonium salt produced undergoes a coupling reaction with 2-naphthol.



(a) Phenylamine, used in Reaction 1, is labelled with the following hazard symbols.

State the meaning of each symbol.

(2)

(b) Practical instructions for the preparation of 1-phenylazo-2-naphthol are shown.

Step 1 Phenylamine, dissolved in hydrochloric acid in a beaker, is placed in an ice bath.

Step 2 When the temperature is between 0°C and 5°C , a cooled aqueous solution of sodium nitrite is added drop-by-drop.

Step 3 A solution of 2-naphthol dissolved in aqueous sodium hydroxide in a boiling tube is placed in an ice bath.

Step 4 The mixture from **Step 3** is then added very slowly to the solution from **Step 2** containing the diazonium salt. The small, red crystals of the product, 1-phenylazo-2-naphthol, form over several minutes.

Step 5 The crystals are filtered off under reduced pressure, washed and recrystallised from ethanoic acid. A data book gives the melting temperature of 1-phenylazo-2-naphthol as 131°C .

- (i) Explain why the temperature for Reaction **1** must be kept between 0°C and 5°C .

(2)

- (ii) Draw a labelled diagram of the apparatus used to filter off the crystals under reduced pressure in **Step 5**.

(2)



- (iii) Outline how the melting temperature of the crystals may be measured to ensure that an accurate value is obtained.

(3)

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Azo dyes were developed during the nineteenth century but workers in the industry showed high rates of bladder cancer and the production of some dyes was discontinued.

The introduction of a sulfonic acid group into the dye used for black ink for ink-jet printers has significantly reduced the toxic effects.

The formula for a sulfonic acid side group is shown.



Suggest how the change in solubility caused by the introduction of the sulfonic acid group reduces the toxic effects of the dye.

(1)

.....

.....

.....

(Total for Question 4 = 10 marks)

TOTAL FOR PAPER = 50 MARKS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



The Periodic Table of Elements

1.0

H

hydrogen

1

10.8

B

boron

5

12.0

C

carbon

6

14.0

N

nitrogen

7

16.0

O

oxygen

8

19.0

F

fluorine

9

20.2

Ne

neon

10

27.0

Al

aluminium

13

28.1

Si

silicon

14

31.0

P

phosphorus

15

32.1

S

sulfur

16

35.5

Cl

chlorine

17

39.9

Ar

argon

18

69.7

Ga

gallium

31

72.6

Ge

germanium

32

74.9

As

arsenic

33

79.0

Se

selenium

34

79.9

Br

bromine

35

83.8

Kr

krypton

36

114.8

In

indium

49

118.7

Sn

tin

50

121.8

Sb

antimony

51

127.6

Te

tellurium

52

126.9

I

iodine

53

131.3

Xe

xenon

54

204.4

Tl

thallium

81

207.2

Pb

lead

82

209.0

Bi

bismuth

83

209.0

Po

polonium

84

210

At

astatine

85

222

Rn

radon

86

4.0

He

helium

2

6.9

Li

lithium

3

9.0

Be

beryllium

4

23.0

Na

sodium

11

24.3

Mg

magnesium

12

39.1

K

potassium

19

40.1

Ca

calcium

20

85.5

Rb

rubidium

37

87.6

Sr

strontium

38

132.9

Cs

caesium

55

137.3

Ba

barium

56

223

Fr

francium

87

226

Ra

radium

88

227

Ac*

actinium

89

261

Rf

rutherfordium

104

262

Db

dubnium

105

266

Sg

seaborgium

106

264

Bh

bohrium

107

277

Hs

hassium

108

271

Ds

darmstadtium

110

272

Rg

roentgenium

111

140

Ce

cerium

58

141

Pr

praseodymium

59

144

Nd

neodymium

60

147

Pm

promethium

61

150

Sm

samarium

62

152

Eu

europium

63

157

Gd

gadolinium

64

159

Tb

terbium

65

163

Dy

dysprosium

66

165

Ho

holmium

67

167

Er

erbium

68

169

Tm

thulium

69

173

Yb

ytterbium

70

175

Lu

lutetium

71

232

Th

thorium

90

231

Pa

protactinium

91

238

U

uranium

92

237

Np

neptunium

93

242

Pu

plutonium

94

243

Am

americium

95

247

Cm

curium

96

245

Bk

berkelium

97

251

Cf

californium

98

254

Es

einsteinium

99

253

Fm

fermium

100

256

Md

mendelevium

101

254

No

nobelium

102

257

Lr

lawrencium

103

* Lanthanide series

* Actinide series

Elements with atomic numbers 112-116 have been reported but not fully authenticated

