OXFORD

INTERNATIONAL AQA EXAMINATIONS

Please write clearly in block capitals.

Centre number | | | | | | Candidate number | | | | |

Surname _____

Forename(s) _____

Candidate signature _____

INTERNATIONAL A-LEVEL

CHEMISTRY (9620)

Unit 4: Organic 2 and Physical 2

Tuesday 22 January 2019 07:00 GMT Time allowed: 1 hour 30 minutes

Materials

For this paper you must have:
- the Periodic Table/Data Sheet, provided as an Insert
- a ruler with millimetre measurements
- a scientific calculator, which you are expected to use where appropriate.

Instructions
- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information
- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
TOTAL	

Answer **all** questions in the spaces provided.

Do not write outside the box

0 1 The rate of reaction between solutions of **A** and **B** is investigated.

A + **B** → products

Three experiments are completed at a given temperature. The results of these experiments are shown in **Table 1**.

Table 1

	Initial concentration of A / mol dm^{-3}	**Initial concentration of B / mol dm^{-3}**	**Initial rate / mol dm^{-3} s^{-1}**
Experiment 1	2.40×10^{-3}	2.80×10^{-3}	1.60×10^{-4}
Experiment 2	3.60×10^{-3}	4.20×10^{-3}	3.60×10^{-4}
Experiment 3	5.40×10^{-3}	4.20×10^{-3}	8.10×10^{-4}

0 1 . 1 Use the data in **Table 1** to deduce the order of reaction with respect to **A** and the order of reaction with respect to **B**.

Place **one** tick (✓) in the box which shows the order with respect to **A** and the order with respect to **B**.

[2 marks]

Order with respect to **A**

Zero ☐

First ☐

Second ☐

Order with respect to **B**

Zero ☐

First ☐

Second ☐

Do not write outside the box

0 1 . 2 The reaction between compounds **C** and **D** is studied at a given temperature.
The rate equation for the reaction is found to be

$$rate = k\,[\mathbf{C}]^2[\mathbf{D}]$$

When the initial concentration of **C** is 8.55×10^{-2} mol dm^{-3} and the initial concentration of **D** is 2.80×10^{-2} mol dm^{-3}, the initial rate of reaction is 9.43×10^{-4} mol dm^{-3} s^{-1}

Calculate the value of the rate constant at this temperature and deduce its units.

[3 marks]

Rate constant _____ Units _____

5

Turn over for the next question

Do not write outside the box

0 2 Iodide ions react with hydrogen peroxide to form iodine as shown.

$$H_2O_2 + 2H^+ + 2I^- \rightarrow I_2 + 2H_2O$$

A student reacted iodide ions with hydrogen peroxide at different temperatures to determine the activation energy (E_a) for this reaction.

The student measured the time taken for a set amount of iodine to be formed at different temperatures. The results from the experiment are shown in **Table 2**.

An equation that relates activation energy (E_a) to the time taken (*time*) and temperature (T), where C is a constant, is shown.

$$\ln\left(\frac{1}{time}\right) = -\frac{E_a}{RT} + \ln C$$

Table 2

Experiment	Temperature / K	Time / s	$\frac{1}{time}$ / s^{-1}	$\ln\left(\frac{1}{time}\right)$	$\frac{1}{T}$ / K^{-1}
1	306	350	2.86×10^{-3}	–5.86	3.27×10^{-3}
2	316	102	9.80×10^{-3}	–4.62	3.16×10^{-3}
3	326	35	2.86×10^{-2}	–3.56	3.07×10^{-3}
4	336	12	8.33×10^{-2}	–2.48	2.98×10^{-3}

0 2 . 1 On **Figure 1**, plot a graph of $\ln\left(\frac{1}{time}\right)$ against $\frac{1}{T}$

You should plot $\ln\left(\frac{1}{time}\right)$ on the *y*-axis and $\frac{1}{T}$ on the *x*-axis.

Draw a line of best fit through your points.


Use the gradient of your line to calculate the activation energy, in kJ mol^{-1}, for this reaction.

The gas constant, R = 8.31 J K^{-1} mol^{-1}

[5 marks]

Do not write outside the box

Figure 1



Activation Energy _____ kJ mol^{-1}

0 2 . 2 Suggest why the student did not repeat the experiment at temperatures higher than 336 K

[1 mark]

_____

_____

6

Turn over ►

Do not write outside the box

0 3 This question is about carboxylic acids and esters.

0 3 . 1 Butanoic acid is a weak acid.

Write an expression for the acid dissociation constant (K_a) for butanoic acid.

[1 mark]

K_a

0 3 . 2 At 298 K for butanoic acid, $K_a = 1.51 \times 10^{-5}$ mol dm^{-3}

Calculate the pH of a 0.150 mol dm^{-3} solution of butanoic acid at 298 K

Give your answer to **two** decimal places.

[3 marks]

pH _____

0 3 . 3 Write an equation to show the reaction of butanoic acid with propan-1-ol.

Identify a catalyst for this reaction.

[2 marks]


Equation

_____

Catalyst _____

Do not write outside the box

0 3 . 4 The structure of a tri-ester is shown.



Give the formula of each of the products formed when this tri-ester is hydrolysed in acid conditions.

[2 marks]

Product 1

Product 2

8

Turn over for the next question

Do not write outside the box

0 4 Propylamine can be formed in different reactions either from propanenitrile or from 1-bromopropane.

0 4 . 1 State the reagent and condition, other than temperature, for the formation of propylamine from each of these starting materials.

[3 marks]

From propanenitrile

Reagent _____

Condition _____

From 1-bromopropane

Reagent _____

Condition _____

0 4 . 2 N-methylethylamine is a secondary amine and an isomer of propylamine.

Suggest why the secondary amine is a stronger base than propylamine.

[2 marks]

_____

_____

_____

_____

5

Do not write outside the box

Turn over for the next question

DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED

Do not write outside the box


0 5 Simple test-tube reactions can be used to distinguish the following pairs of isomers.

Give a reagent, or combination of reagents, that can be added to the compounds in each pair to distinguish between them.


State what is observed in each case.

0 5 . 1

R



S



[3 marks]


Reagent _____

Observation with **R** _____


Observation with **S** _____

0 5 . 2

T



U



[3 marks]

Reagent _____


Observation with **T** _____

Observation with **U** _____


Do not write outside the box

0 5 . 3

V



W



[3 marks]

Reagent _____

Observation with **V** _____

Observation with **W** _____


9

Turn over for the next question


Do not write outside the box

0 6 Sections of two polymers, **E** and **F**, are shown.

E



F



0 6 . 1 Polymer **E** is made from two different alkene monomers.

Draw the skeletal formula of each alkene monomer.

[2 marks]

Alkene monomer 1 **Alkene monomer 2**

Do not write outside the box

0 6 . 2 Polymer **F** is a polyester. This polyester is produced by reacting a diol with a dicarboxylic acid.

Give the structure of the diol and the structure of the dicarboxylic acid used to produce this polyester.

[2 marks]

Diol

Dicarboxylic acid

0 6 . 3 Suggest an environmental advantage of polymer **F** compared with polymer **E**.

Explain your answer.

[3 marks]

Advantage _____

Explanation _____

_____

_____


_____

7

Turn over ►

Do not write outside the box

0 7 The structures of three amino acids are shown.



Serine **Cysteine** **Aspartic acid**

0 7 . 1 Use IUPAC rules to deduce the name for serine.

[1 mark]

0 7 . 2 Draw the structure of the zwitterion formed by cysteine.

[1 mark]

0 7 . 3 Draw the structure of the species formed by aspartic acid at high pH.

[1 mark]

Do not write outside the box

0 7 . 4 Draw the structure of each of the two dipeptides formed when cysteine and serine react together.

[2 marks]

Dipeptide 1

Dipeptide 2

0 7 . 5 Draw the structure of the species formed when cysteine reacts with an excess of bromomethane.

[1 mark]

Do not write outside the box

0 7 . 6 Draw the structure of the species formed when serine reacts with an excess of ethanoyl chloride.

[2 marks]

0 7 . 7 A mixture of amino acids can be separated using thin-layer chromatography.

Describe how amino acids can be made visible on the chromatogram.

Explain why different amino acids have different R_f values.

[3 marks]

How amino acids are made visible _____

_____

Explanation _____

_____

_____

_____

_____

_____

Do not write outside the box

0 7 . 8 **Table 3** shows different types of bonding in proteins.

Complete **Table 3** by placing a tick (✓) in the correct column to show which level of protein structure (primary, secondary or tertiary) contains each type of bonding.

[2 marks]

Table 3

Type of bonding	Primary	Secondary	Tertiary
Hydrogen bonds in an α-helix			
Peptide links between amino acids			
Hydrogen bonds between $-CH_2OH$ groups in serine amino acids			
Disulfide bonds between cysteine amino acids			

13


Turn over for the next question

Do not write outside the box

0 8 Compound **G** has the molecular formula $C_4H_8O_2$

The infrared spectrum of **G** is shown in **Figure 2**.

Figure 2



0 8 . 1 Use **Table A** on the Data Sheet to identify the bond causing the absorption labelled **I** and the bond causing the absorption labelled **II**.

[2 marks]

Bond **I** _____

Bond **II** _____

0 8 . 2 Proton (^1H) NMR spectra are recorded using solutions that contain the non-toxic and inert standard tetramethylsilane (TMS).

Give **two** other reasons why TMS is a suitable standard.

[2 marks]

Reason 1 _____

_____

Reason 2 _____

_____

Do not write outside the box

Question 8 continues on the next page

DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED

Do not write outside the box

0 8 . 3 Identify a suitable solvent to use when recording a ^{1}H NMR spectrum.

Give a reason for your choice.

[2 marks]

Solvent _____

Reason _____

The ^{1}H NMR spectrum of **G** shows four peaks. **Table 4** gives chemical shift (δ) values for the four peaks, their splitting patterns and integration values.

Table 4

δ / ppm	2.20	2.60	3.40	3.84
Splitting pattern	singlet	triplet	singlet	triplet
Integration value	0.75	0.50	0.25	0.50

Use **Figure 2**, **Table 4** and **Table B** of your Data Sheet to help you answer questions **08.4**, **08.5** and **08.6**.

0 8 . 4 Draw the part of the structure of **G** that can be deduced from the peak at δ = 2.20 ppm

[1 mark]

0 8 . 5 Draw the part of the structure of **G** that can be deduced from the peak at δ = 3.40 ppm

[1 mark]

Do not write outside the box

0 8 . 6 Draw the part of the structure of **G** that can be deduced from the peaks at δ = 2.60 ppm and δ = 3.84 ppm

[2 marks]

0 8 . 7 Deduce the structure of **G**.

[1 mark]


11

Turn over for the next question

Do not write outside the box

0 9 1-Phenylpropene can be prepared as shown in **Figure 3**.

Figure 3



0 9 . 1 **Step 1** involves an attack on benzene by the electrophile $[CH_3CH_2CO]^+$


Write an equation for a reaction that forms this electrophile.

Complete the mechanism for the reaction of this electrophile with benzene.

[3 marks]

Equation _____

Mechanism



Do not write outside the box

0 9 . 2 **Step 2** involves the reaction of **L** with $NaBH_4$

Name and outline the mechanism for this reaction.

[5 marks]

Name of mechanism _____

Mechanism

Question 9 continues on page 25

Do not write outside the box

There are no questions printed on this page

DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED


Do not write outside the box

0 9 . 3 1-Phenylpropene can be formed by reacting **M** with concentrated sulfuric acid. Part of the mechanism is shown in **Figure 4**.

Draw **two** curly arrows on **Figure 4** to complete the mechanism.
You do **not** need to draw the final products.

[2 marks]

Figure 4




Question 9 continues on the next page

Do not write outside the box

Figure 3 is repeated here.

Figure 3



0 9 . 4 **M** exists as two enantiomers.

State the meaning of the term enantiomers.

[1 mark]

0 9 . 5 Describe how the two enantiomers could be distinguished.

[2 marks]

Do not write outside the box

0 9 . 6 The overall percentage yield in the preparation of 1-phenylpropene from benzene is 68.0%

Calculate the minimum mass of benzene needed to produce 1.68 g of 1-phenylpropene.

[3 marks]

Mass of benzene _____ g

16

END OF QUESTIONS

Do not write outside the box

There are no questions printed on this page

DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED

Do not write outside the box

Question number	**Additional page, if required.** **Write the question numbers in the left-hand margin.**

Do not write outside the box

Question number	**Additional page, if required.** **Write the question numbers in the left-hand margin.**

Do not write outside the box

Question number	**Additional page, if required.** **Write the question numbers in the left-hand margin.**

Do not write outside the box

Question number	**Additional page, if required.** **Write the question numbers in the left-hand margin.**

Copyright information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third-party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2019 AQA and its licensors. All rights reserved.