OXFORD

INTERNATIONAL AQA EXAMINATIONS

Please write clearly in block capitals.

Centre number | | | | | | Candidate number | | | | |

Surname _____

Forename(s) _____

Candidate signature _____

INTERNATIONAL A-LEVEL
FURTHER MATHEMATICS

(9665/FM03) Unit FP2 − Pure Maths

Friday 21 June 2019 07:00 GMT Time allowed: 2 hours 30 minutes

Materials
- For this paper you must have the Oxford International AQA booklet of formulae and statistical tables (enclosed).
- You may use a graphics calculator.

Instructions
- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information
- The marks for questions are shown in brackets.
- The maximum mark for this paper is 120.

Advice
- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- Show all necessary working; otherwise marks may be lost.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
TOTAL	

Do not write outside the box

Answer **all** questions in the spaces provided.

1 (a) Show that

$$\frac{1}{2r+1} - \frac{1}{2r+3} = \frac{k}{(2r+1)(2r+3)}$$

where k is a constant.

[2 marks]

Do not write outside the box

1 (b) Hence use the method of differences to show that

$$\sum_{r=1}^{n} \frac{1}{(2r+1)(2r+3)} = \frac{n}{p(2n+3)}$$

where p is an integer.

[4 marks]

6

Turn over for the next question

Do not write outside the box

2 (a) Use the definition of $\cosh x$ in terms of e^{x} and e^{-x} to show that

$$2\cosh^2 x - 1 = \cosh 2x$$

[3 marks]

Do not write outside the box

2 (b) A curve has equation

$$y = 3\sinh 2x - 5\sinh x + 4x$$

Using the result in part **(a)**, prove that this curve has no stationary points.

[6 marks]

9

Do not write outside the box

3 The roots of the cubic equation

$$3z^3 + 9z + r = 0$$

where r is real, are α, β and γ.

3 (a) (i) Write down the value of $\alpha\beta + \beta\gamma + \gamma\alpha$.

[1 mark]

$\alpha\beta + \beta\gamma + \gamma\alpha =$ _____

3 (a) (ii) Hence show that $\alpha^2 + \beta^2 + \gamma^2 = -6$

[3 marks]

Do not write outside the box

3 (a) (iii) Hence explain why the cubic equation must have two non-real roots and one real root.

[2 marks]

3 (b) (i) Given that $\alpha = 1 + \sqrt{6}\,\mathrm{i}$, find the value of $\alpha\beta\gamma$.

[3 marks]

$\alpha\beta\gamma =$ _____

3 (b) (ii) Hence write down the value of r.

[1 mark]

$r =$ _____

10

Do not write outside the box

4 Three planes have equations

$$x + 3y + cz = c + 4$$
$$x + 2y + 3z = 6$$
$$x + y + z = d$$

where c and d are constants.

The three planes do not intersect at a unique single point.

4 (a) Show that $c = 5$

[2 marks]

Do not write outside the box

4 (b) In the case where the three planes also share a common line of intersection, determine the value of d.

[5 marks]

$d =$ _____

7

Turn over for the next question

Do not write outside the box

5 (a) Given that

$$f(k) = 2^{k+2} + 3^{2k+1}$$

show that

$$f(k+1) - 2f(k) = a \times 3^{2k+1}$$

where a is an integer.

[3 marks]

Do not write outside the box

5 (b) Hence prove by induction that $2^{n+2} + 3^{2n+1}$ is a multiple of 7 for all integers $n \geq 1$

[4 marks]

7

Turn over ►

Do not write outside the box

6 It is given that y satisfies the differential equation

$$\frac{\mathrm{d}^2y}{\mathrm{d}x^2} + \frac{\mathrm{d}y}{\mathrm{d}x} - 6y = 20\,\mathrm{e}^{2x} + 18$$

6 (a) Find the values of the constants p and q for which $p + qx\mathrm{e}^{2x}$ is a particular integral of this differential equation.

[5 marks]

$p =$ _____ $q =$ _____

Do not write outside the box

6 (b) Hence solve the differential equation, expressing y in terms of x, given that $y = 5$ when $x = 0$ and that $\frac{dy}{dx} \to 0$ as $x \to -\infty$

[7 marks]

Answer _____

12

Do not write outside the box

7 The matrix $\mathbf{M} = \begin{bmatrix} 1 & -1 & 2 \\ 0 & 5 & 7 \\ k & 1 & 1 \end{bmatrix}$, where k is a positive integer.

An eigenvector of $\mathbf{M}$ is $\begin{bmatrix} -1 \\ 7 \\ 1 \end{bmatrix}$ and its corresponding eigenvalue is λ_1

7 (a) Find the value of λ_1 and the value of k.

[4 marks]

$\lambda_1 =$ _____ $k =$ _____

7 (b) Show that -2 is the least eigenvalue of $\mathbf{M}$.

[4 marks]

Do not write outside the box

7 (c) Find an eigenvector corresponding to the eigenvalue -2

[3 marks]

Answer _____

7 (d) The transformation T has matrix $\mathbf{M}$.

Write down the Cartesian equations for any one of the invariant lines of T.

[1 mark]

Answer _____

12

Do not write outside the box

8 (a) Use integration to show that $\frac{x}{x+1}$ is an integrating factor for the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} + \frac{1}{x(x+1)}y = 2x + 3\,, \quad x > 0$$

[4 marks]

Do not write outside the box

8 (b) Hence find the general solution of this differential equation.

[5 marks]

Answer

9

Do not write outside the box

9 Plane Π_1 has vector equation $\mathbf{r}.\begin{bmatrix} 2 \\ -1 \\ 2 \end{bmatrix} = 3$

Plane Π_2 has vector equation $\mathbf{r}.\begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix} = 4$

9 (a) Calculate the acute angle between the planes Π_1 and Π_2 giving your answer to the nearest 0.1°

[4 marks]

Answer _____

Do not write outside the box

9 (b) Find a vector equation for the line of intersection of Π_1 and Π_2

[5 marks]

Answer _____

9

Do not write outside the box

10 A curve C is given parametrically by the equations

$$x = t - \sin t \ , \qquad y = 2\sin^2\left(\frac{t}{2}\right)$$

10 (a) Show that $\left(\frac{\mathrm{d}x}{\mathrm{d}t}\right)^2 + \left(\frac{\mathrm{d}y}{\mathrm{d}t}\right)^2 = 4\sin^2\left(\frac{t}{2}\right)$.

[4 marks]

Do not write outside the box

10 (b) The arc of C between the points where $t = \frac{\pi}{3}$ and $t = \frac{\pi}{2}$ is rotated through 2π radians about the x-axis.

Find the area of the surface generated, giving your answer in the form $\frac{2\pi}{3}\left(p\sqrt{3} + q\sqrt{2}\right)$, where p and q are integers.

[6 marks]

Answer _____

10

Turn over ►

Do not write outside the box

11 Given that $z = \cos\theta + \mathrm{i}\sin\theta$:

11 (a) (i) use de Moivre's theorem to show that $z^n + \frac{1}{z^n} = 2\cos n\theta$.

[3 marks]

11 (a) (ii) write down an expression for $z - \frac{1}{z}$ in terms of $\sin\theta$.

[1 mark]

Answer _____

Do not write outside the box

11 (b) Hence express $64\sin^6\theta$ in the form

$$20 + a\cos 2\theta + b\cos 4\theta + c\cos 6\theta$$

where a, b and c are integers.

[5 marks]

Answer _____


Question 11 continues on the next page

Do not write outside the box

11 (c) A leaf lies flat on a thin circular disc.

The diagram shows a curve C_1 which models the leaf and a circle C_2 which models the disc.

The pole O and the initial line are also shown.



The polar equation of the curve C_1 is $r = 2\sin^3\theta$, $0 \leq \theta \leq \pi$.

The polar equation of the circle C_2 is $r = 2\sin\theta$, $0 \leq \theta \leq \pi$.

Using your answer to part **(b)**, find what percentage of the area of the circular disc is **not** covered by the leaf.

[5 marks]

Do not write outside the box

Answer _____

14

Do not write outside the box

12 (a) Write down the expansion of $\sin 2x$ in ascending powers of x up to and including the term in x^5

[1 mark]

Answer _____

12 (b) (i) Given that

$$\ln y = \tan^{-1} x$$

prove that

$$(2x-1)\left(\frac{dy}{dx}\right)^2 + y\frac{d^2y}{dx^2} = 0$$

[5 marks]

Do not write outside the box

12 (b) (ii) Hence, given that the first five terms in the Maclaurin series expansion in ascending powers of x of $e^{\tan^{-1}x}$ are

$$1 + x + \frac{x^2}{2} + px^3 + qx^4$$

show that $p = -\frac{1}{6}$ and find the value of q.

[6 marks]

Do not write outside the box

$q =$ _____

12 (c) Hence show that $\lim_{x \to 0} \left[\frac{e^{\tan^{-1} x} - e^{x}}{2x - \sin 2x} \right]$ exists and find its value.

[3 marks]

Answer _____

END OF QUESTIONS

15

Do not write outside the box

There are no questions printed on this page

DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED

Do not write outside the box

Question number	**Additional page, if required.** **Write the question numbers in the left-hand margin.**

Do not write outside the box

Question number	**Additional page, if required.** **Write the question numbers in the left-hand margin.**

Do not write outside the box

Question number	**Additional page, if required.** **Write the question numbers in the left-hand margin.**

Copyright information

For confidentiality purposes, acknowledgements of third-party copyright material are published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.oxfordaqaexams.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and Oxford International AQA Examinations will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2019 Oxford International AQA Examinations and its licensors. All rights reserved.